

New study shows molecular details of how organic aerosol helps heat up and color haze over megacities

September 29 2015



Scientists determined that high concentrations of nitrogen oxides influence the creation of the brown haze that hangs over megacities, providing new insights that can create higher accuracy climate and atmospheric models. Credit: Alex Laskin

Sitting on an airplane flying into nearly every major city on a sunny day, passengers can see a lingering brown haze. This haze event is linked to

climate issues, as the thousands of chemicals involved act as a warming blanket, absorbing sunlight and trapping surface heat. Yet, the chemistry of this haze is not well known. Recently, scientists at Pacific Northwest National Laboratory examined the chemistry of brown carbon, a pernicious set of particles in the haze. They focused on the specks that form around the chemical toluene, a common pollutant. They found that adding a bit of nitrogen oxide, released in combustion engine car exhaust, resulted in particles that trapped some heat. Increasing the nitrogen oxides level, however, resulted in particles that held significantly more heat and caused the mix to turn yellowish brown.

The results are highlighted on the cover of a recent issue of *Physical Chemistry Chemical Physics*.

To deal with today's air quality issues and minimize future problems, scientists need to accurately predict the structure and behavior of the different types of particles involved. This study provides new insights into the evolution and behavior of a less studied type of particle, called "brown carbon." Brown carbon is composed of organic molecules that absorb and retain heat. The results could lead to better predictive models to inform future decisions about energy production and use, and a better understanding of changes in the climate.


The team at PNNL took a make-measure-model approach. Dr. John Shilling and Dr. Jiumeng Liu synthesized the brown carbon particles by mixing the common anthropogenic pollutant toluene with nitrogen oxides inside PNNL's environmental chamber. They exposed the resulting gas to light, photo-oxidizing the mixture to mimic the natural aging process that occurs in the atmosphere. The result was particles with characteristic brown carbon properties.

They passed the brown carbon material to Dr. Alex Laskin, Dr. Julia Laskin, and Dr. Peng Lin. Using analytical capabilities in EMSL,

including high-resolution mass spectrometry, they investigated the molecular structures and optical properties of chemical components of this material. With the experimental results in hand, Dr. Shawn Kathmann ran a series of detailed calculations. He identified the excited electron states in 15 chromophores, or light-absorbing structures, and provided theoretical chemistry insights on the mechanisms by which these molecules take in the light. This result starts to build the roster of players that form the haze observed over cities and lets scientists better predict the properties of other particles.

PCCP

Physical Chemistry Chemical Physics
www.rsc.org/pccp



ISSN 1463-9076




PAPER
Alexander Laskin et al.
Molecular characterization of brown carbon (BrC) chromophores in
secondary organic aerosol generated from photo-oxidation of toluene

The research showing how a mixture of nitrogen oxides and toluene increase light-absorbing properties of brown carbon particles appeared on the cover of *Physical Chemistry Chemical Physics*. Credit: Pacific Northwest National

Laboratory and the Royal Society of Chemistry in PCCP 17:23312.

The team is doing further studies on the properties of [brown carbon](#) to inform computational models to assess energy choices and their influence on the climate.



Testing candidate molecular structures that resulted from the mass spectrometry studies of brown carbon material, the team found that three molecules absorbed light in the visible part of the spectrum (on the bottom row, the 1st, 2nd, and 4th from the left). The atom colors are cyan (carbon), red (oxygen), blue (nitrogen), and white (hydrogen).

More information: "Molecular Characterization of Brown Carbon (BrC) Chromophores in Secondary Organic Aerosol Generated from Photo-Oxidation of Toluene." *Physical Chemistry Chemical Physics* 17:23312. [DOI: 10.1039/c5cp02563j](https://doi.org/10.1039/c5cp02563j)

Provided by Pacific Northwest National Laboratory

Citation: New study shows molecular details of how organic aerosol helps heat up and color haze over megacities (2015, September 29) retrieved 11 July 2024 from <https://phys.org/news/2015-09-ew-molecular-aerosol-haze-megacities.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.