

Faster resistance analysis for patients with blood poisoning

September 25 2015



A new technique for analyzing antibiotic resistance. Credit: Volker Lannert, Fraunhofer FIT

When treating cases of blood poisoning, doctors resort immediately to broad-spectrum antibiotics. The problem is that in many cases the



bacteria are resistant to the medicine. Analyzing antibiotic resistance is a time-consuming process, and for many patients the results come too late. Now a new technique that supplies results in just nine hours is to be presented at the Biotechnica trade show in Hanover from October 6-8 (Hall 9, booth C34).

For patients with <u>blood poisoning</u>, also known as septicemia, every second counts. Doctors who suspect a patient has sepsis start them on <u>broad-spectrum antibiotics</u> right away, but the <u>antibiotics</u> don't always have the desired effect – for instance if the <u>bacteria</u> are resistant to the medicines used. Identifying the pathogens in the lab and investigating their potential resistance routinely takes between 60 and 100 hours. This is time the patient simply doesn't have – most cases end fatally within around 48 hours. Blood poisoning accounts for 60,000 deaths a year in Germany alone.

Test results in nine hours

In future, this analysis could take much less time – saving many patients' lives. Once doctors know whether the bacteria are resistant to certain substances, they can treat the patient with a targeted antibiotic that reliably kills off the pathogen. This is possible thanks to a technology developed by the researchers at the Fraunhofer Institutes for Applied Information Technology FIT and for Laser Technology ILT in collaboration with the Uniklinikum Aachen and numerous industry partners. "Our testing method yields results in just nine hours," says Professor Harald Mathis, department head at Fraunhofer FIT.

Which antibiotic?

So how are researchers now able to analyze the bacteria in a blood sample up to ten times faster than before? "We've developed a



miniaturized system with a patented optical design," reveals Mathis. The first step is to mark the pathogens indicative of septicemia, so that they glow when exposed to laser light. This then allows the researchers to assess the amount of bacteria present in the blood. In the next stage of the process, the bacteria are separated from the blood and channeled into a series of miniaturized dishes. Each contains a culture medium that includes a specific antibiotic. A second optical system complete with the necessary analysis software observes and precisely documents how the bacteria develop. Then comes the key step: algorithms analyze the pictures taken of the bacteria and extrapolate the growth curve, meaning the researchers can see within hours whether the respective medicine is working or whether the bacteria are resistant to it and spreading rapidly. Essentially, the growth monitor software is able to calculate and predict how bacteria will develop over time.

It does so by analyzing both the extent of the bacterial growth – which provides a one-to-one indication of the number of bacteria present – and the ratio of living to dead bacteria. In short: This tells researchers which antibiotic will be most effective in killing off the bacteria – and help the patient the most.

Provided by Fraunhofer-Gesellschaft

Citation: Faster resistance analysis for patients with blood poisoning (2015, September 25) retrieved 30 April 2024 from https://phys.org/news/2015-09-faster-resistance-analysis-patients-blood.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.