

# Study provides insights into the mechanisms of fine-tuning of wheat to diverse environments

September 2 2015

---



A recent *Proceedings of the National Academy of Sciences* study identifies the last major vernalization gene in wheat. The gene, VRN-D4, is found in wheat lines primarily grown in South Asia, particularly northeastern Pakistan and northwestern India. The gene adapted the wheat varieties grown in the region to the local environment. Credit: Kansas State University

A Kansas State University wheat geneticist is part of a breakthrough study that identifies one of the wheat genes that controls response to low temperature exposure, a process called vernalization. Natural variation in vernalization genes defines when the plant begins to flower and is critical for adaptation to different environments.

Researchers anticipate this will help wheat breeders design wheat varieties that can adapt and thrive in changing environments around the world.

Eduard Akhunov, associate professor in the plant pathology department, collaborated with Jorge Dubcovsky at the University of California, Davis on this study. It appears in the *Proceedings of the National Academy of Science*, or *PNAS*, article "Identification of the VERNALIZATION 4 gene reveals the origin of spring growth habit in ancient wheats from South Asia."

"The vernalization [genes](#) can sense for how long the plant was subjected to low temperature treatment and trigger flowering when the time is right," Akhunov said. "Wheat adapted to different environmental conditions by using different forms of these vernalization genes—the first three of which had been identified by Dr. Dubcovsky's group prior to this study. This is the last major vernalization gene to complete the story."

In the study, researchers investigated a gene called VRN-D4 that is found in [wheat lines](#) primarily grown in South Asia, particularly northeastern Pakistan and northwestern India. They showed that the gene likely originated by duplication of a major vernalization gene VRN-A1.

Researchers analyzed the genetic diversity in the global wheat populations to test if the VRN-D4 gene helped the wheat varieties grown in South Asia adapt to the region's climate and environment. They used a

variety of population genetics and bioinformatics approaches to analyze genetic diversity along the chromosomes in multiple wheat varieties around the world.

Researchers looked for regions of each plant's genome that showed unusual patterns of variation consistent with past selective breeding acting on the VRN-D4 gene. The team observed that while the VRN-D4 gene is present in wheat varieties grown in different parts of the world, it is most frequent in South Asia.

The researchers concluded it is likely that ancient farmers favored the VRN-D4 variant of the vernalization gene in this region because of its high adaptive value in the local environment.

"A wheat line's adaptation to its local environment is probably one of the most important properties of the plant for wheat breeders because it directly influences crop production," Akhunov said. "If you grow lines of [wheat](#) that are not adapted to a specific environment, you will not get a very high yield."

Plant scientists have only recently begun to understand how the vernalization response in plants works, Akhunov said. The latest findings offer new focus points into how these genes and others may be modified to fine-tune a [wheat variety](#) for a particular environment, which will result in less crop and food loss due to changing environment.

**More information:** Identification of the VERNALIZATION 4 gene reveals the origin of spring growth habit in ancient wheats from South Asia, Nestor Kippes, [DOI: 10.1073/pnas.1514883112](https://doi.org/10.1073/pnas.1514883112)

Provided by Kansas State University

Citation: Study provides insights into the mechanisms of fine-tuning of wheat to diverse environments (2015, September 2) retrieved 20 March 2024 from <https://phys.org/news/2015-09-insights-mechanisms-fine-tuning-wheat-diverse.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.