

Study uncovers new approaches for sorghum breeders

September 10 2015



A recent study on plant height in sorghum will likely be applicable to other economically important traits, such as crop yield. Credit: Kansas State University

A study on a sorghum population at Kansas State University has helped researchers better understand why a crop hybrid often performs better

than either of its parent lines, known as heterosis.

The findings are specific to plant height in [sorghum](#), but a Kansas State University agronomist says they will likely be applicable to other economically important traits, such as crop yield.

The research, "Dissecting repulsion linkage in the dwarfing gene Dw3 region for sorghum plant height provides insights into heterosis," was published Sept. 8 in the *Proceedings of the National Academy of Sciences*.

"Heterosis is a complex genetic phenomena that has been theorized in many different ways," said Tesfaye Tesso, a Kansas State University associate professor of agronomy. "This study adds another dimension to that complexity by pinpointing the impact of repulsion phase linkage on heterosis."

The repulsion phase linkage refers to the link between dominant alleles—forms of the gene responsible for hereditary characteristics—of one or more genes, and the recessive alleles of another.

In repulsion phase linkage, one parent contributes the dominant allele of one gene and the second parent—often more than one—provides the dominant allele of other genes. In the current study, researchers successfully laid out how such linkage between genes associated with plant height can affect heterosis for a given trait.

"Plant height is a relatively simple trait and is easier to track in segregating a population," Tesso said. "Other economically important traits such as yield are more complex, but like plant height, heterosis has a significant contribution to yield. If we know as much information about heterosis for yield as we do for plant height, it will increase our ability to select parental lines that are most compatible and give

maximum heterosis."

Tesso managed the sorghum population used in this study in Kansas State University research fields. The population is a recombinant inbred line, which allows researchers to produce large amounts of seed for replicated trials.

Kansas State University maintains the sorghum germplasm that was used for this research. Jianming Yu, a former Kansas State University professor who is now at Iowa State, led the study. The work involves experts in plant breeding, genetics and physiology.

Tesso said that plant breeders can immediately use these findings to influence [plant height](#) in new sorghum lines. Eventually, the results will help to improve many other desirable traits in sorghum.

Sorghum is an important crop around the world, especially in many African countries where it's estimated that hundreds of millions of people rely on the crop as a food source.

The United States is the world's top exporter of [grain sorghum](#), accounting for almost 75 percent of global trade, according to the U.S. Grains Council. China, which primarily uses the crop for livestock feed, is the top importer of U.S. sorghum.

More information: Dissecting repulsion linkage in the dwarfing gene Dw3 region for sorghum plant height provides insights into heterosis, *PNAS*, [DOI: 10.1073/pnas.1509229112](https://doi.org/10.1073/pnas.1509229112)

Provided by Kansas State University

Citation: Study uncovers new approaches for sorghum breeders (2015, September 10) retrieved 11 July 2024 from <https://phys.org/news/2015-09-uncovers-approaches-sorghum-breeders.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.