


Researchers create better algorithm for simulating particles in Fermi Sea

October 27 2015



A single down-spin impurity surrounded by a Fermi sea of up-spin particles in two dimensions.

A North Carolina State University physicist and his German colleagues have created a new, more precise algorithm for simulating particle interactions when a single impurity is introduced into a Fermi sea. The algorithm shows that when these particles interact, the transition from quasiparticle to bound molecule in a polarized two-dimensional system is

smooth. The new method may have implications for understanding the behavior of impurities in a variety of systems.

The Fermi sea describes a collection of weakly interacting identical fermions such as electrons that have been cooled to a very low temperature. No two fermions within the sea have exactly the same quantum state. The ground state of the Fermi sea in this pure form is well understood. However, what happens when an impurity - such as a particle with a different spin - is introduced? How does that one particle affect the system as a whole?

"Let's say that all the particles in the sea are up-spin particles, and we introduce one down-spin particle," says NC State physicist Dean Lee, co-author on a paper describing the work. "Does this new particle form a molecular bond with one of the up-spin particles? How does the system react?"

Lee and his colleagues, lead author Shahin Bour and Ulf-G. Meissner from Bonn University and Hans-Werner Hammer from Darmstadt University, developed a [lattice](#) algorithm called impurity lattice Monte Carlo that samples the possible paths of the impurity in the Fermi sea. Monte Carlo methods are commonly used to simulate quantum mechanical systems. Impurity lattice Monte Carlo differs from other methods in that it treats the impurity particle explicitly, in a completely different manner from the other particles in the system.

According to the lattice results, the transition from single particle to bound molecule is smooth. "Physicists had theorized that there should be a clear critical value, or interaction strength, where the impurity would bond with another particle and become a molecule," Lee says, "but our simulations don't show that. Instead, we find that there's an interesting ambiguous state where the [particles](#) are interacting, but may or may not be a bound molecule. And when the transition does happen, it occurs

smoothly as a function of interaction strength.

"What we're most excited about, though, are the future possibilities. We want to take the lattice into three-dimensional simulations, and introduce an [impurity](#) to a paired superfluid to see what effects that has on the system. We hope that our method can be used to address questions relevant to cold atoms, solid state systems and neutron stars."

The research appears in *Physical Review Letters*.

More information: Shahin Bour et al. Lattice Results for Fermi Polarons in Two Dimensions , *Physical Review Letters* (2015). [DOI: 10.1103/PhysRevLett.115.185301](https://doi.org/10.1103/PhysRevLett.115.185301)

Provided by North Carolina State University

Citation: Researchers create better algorithm for simulating particles in Fermi Sea (2015, October 27) retrieved 26 March 2023 from <https://phys.org/news/2015-10-algorithm-simulating-particles-fermi-sea.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.