


# Climate models used to explain formation of Mars valley networks

October 12 2015, by A'ndrea Elyse Messer

---



Valles Marineris, Mars. Credit: NASA

The extensive valley networks on the surface of Mars were probably created by running water billions of years ago, but the source of that water is unknown. Now, a team of Penn State and NASA researchers is using climate models to predict how greenhouse warming could be the source of the water.

"Everyone is looking for life on Mars, and if Mars was habitable early on as indicated by flowing water, then the chances of there being some sort

of life there now goes up," said Natasha Batalha, graduate student in astronomy and astrophysics, Penn State. "Our work involves using models to best estimate what was happening on the surface of Mars 3.8 billion years ago."

Previous studies analyzing craters support the idea of flowing water on early Mars but still unknown is how surface warming occurred to melt that water and how much water was really there.

"If we compare some of these valleys on Mars to what we know on Earth, for example the Colorado River, then they look to be the same width," said James Kasting, Evan Pugh Professor of Geosciences, Penn State. "We know how much water it took to carve the Grand Canyon based on five to six million years of rainfall, which is estimated to be about three to six million feet (of rainfall)."

In 2014, Kasting's group proposed a climate model showing Mars warmed by a dense atmosphere of greenhouse gases, specifically carbon dioxide and hydrogen. In a recent issue of *Icarus*, they report that using a photochemical model, they determined the possibility that there was a high percentage of hydrogen in the atmosphere. According to Batalha and Kasting, photochemical models simulate different aspects of the atmosphere using mathematics to determine the chemicals and how they react. They also set limits that define what is happening at the surface and in the atmosphere.

"The main focus of our work was how can you, and is it even possible to, get a 5 percent hydrogen atmosphere because 5 percent is a lot. Terrestrial planets don't usually have that big of a hydrogen envelope because it escapes pretty easily," said Batalha.

To get a high percentage of hydrogen in the atmosphere, they found that they need volcanic fluxes caused by plate tectonics. However, the

researchers believe that while there is some evidence that Mars had plate tectonics, this theory remains controversial. Future work will look for evidence that supports their climate model without relying on [plate tectonics](#). One possibility might show that hydrogen escapes more slowly into the atmosphere than currently predicted.

A competing hypothesis, called the [impact hypothesis](#), claims that the high bombardment of asteroids on the surface of Mars led to a temporary warming, causing steam atmospheres that then condensed and produced rainfall that carved the valleys on Mars' surface. However, that would have generated only around 1,500 feet of water, while the Grand Canyon took millions of feet of water to form, according to Kasting.

Data gathered by NASA's Curiosity Rover, launched in 2011 and currently on Mars, suggests prolonged, standing liquid water was present in Gale Crater. According to Batalha, further evidence showing [water](#) stayed on Mars for tens of millions of years would support a warm early Mars while putting the impact hypothesis to rest.

"While Curiosity couldn't directly prove our hypothesis about a high hydrogen atmosphere, if you get millions of years of warmth, then you're driven towards [hydrogen](#) because we haven't been able to figure out any other way to create that warmth. All of the other greenhouse mechanisms fail in our models," added Batalha.

Also working on this project were Shawn D. Domagal-Goldman, research space scientist, NASA, and Ramses Ramirez, a recent Penn State Ph.D in geosciences, now a research associate at Cornell University.

The National Science Foundation, NASA's Exobiology and Astrobiology programs, the Simon's Foundation, and the Carl Sagan Institute supported this work.

Provided by Pennsylvania State University

Citation: Climate models used to explain formation of Mars valley networks (2015, October 12)  
retrieved 23 April 2024 from

<https://phys.org/news/2015-10-climate-formation-mars-valley-networks.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.