

Collaborative research reveals a new view of cell division

October 23 2015

Basic research into the mechanisms of cell division, using eggs and embryos from frogs and starfish, has led researchers to an unexpected discovery about how animal cells control the forces that shape themselves.

During a key point in cytokinesis—the process in which a cell divides its cytoplasm to create two daughter cells—a cell's cortex becomes an excitable medium, the researchers report in a paper placed online ahead of print by the journal *Nature Cell Biology*.

It has long been clear that cytokinesis is highly dynamic, but before it happens an animal cell has no idea where this crucial event will occur, said co-author George von Dassow, a University of Oregon biologist at the Oregon Institute of Marine Biology in Charleston. "After it happens, the cell has literally ripped itself in two," he said. "The contractile machinery that makes this happen is quintessentially ephemeral."

What hasn't been obvious, he said, is how the cell manages to ensure that the entire surface can participate, but, once specified, only one narrow equatorial band does the crucial act. The cell surface enters an excitable state just after chromosomes separate deep in the [cell cytoplasm](#).

At that point, von Dassow said, waves of signaling molecules form. They appear to function like an amplifier, tuning in faint signals from deep in the cell to accurately and precisely delineate the working conditions for contractile proteins and other enzymes to assemble at the right place, in

the right amount and at the right time during [cell division](#), or, as the research team theorized, during other important cell-shape changes.

This "cell-cycle-entrained behavior" in the cortex—a thin layer of organized cytoplasm associated with the cell membrane—is present in vertebrates and invertebrates, the researchers show in their paper.

"Systems as diverse as microbial populations, [cardiac muscle cells](#), the inflammatory response, and even defined chemical mixtures, may exhibit excitability, which has long been an interest to mathematicians," von Dassow said. "High-resolution live-cell imaging combined with mathematical modeling suggests how this dynamic behavior might be used by [cells](#) to accurately and adaptively manage their contractile machinery."

The discovery emerged from work being done under two National Science Foundation grants (MCB-0917887 and MCB-1041200) to the UO's von Dassow and a National Institutes of Health grant (GM52932) to co-author William M. Bement of the University of Wisconsin-Madison.

"We are longtime collaborators, and it just happened that two threads of research accidentally converged on what turned out to be a similar phenomenon in both frog eggs and embryos, and starfish eggs and embryos," von Dassow said.

More information: Activator–inhibitor coupling between Rho signalling and actin assembly make the cell cortex an excitable medium, [nature.com/articles/doi:10.1038/ncb3251](https://doi.org/10.1038/ncb3251)

Provided by University of Oregon

Citation: Collaborative research reveals a new view of cell division (2015, October 23) retrieved 27 April 2024 from <https://phys.org/news/2015-10-collaborative-reveals-view-cell-division.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.