

# Cow-calf grazing practices could determine, mitigate greenhouse gas emissions

October 29 2015, by Kay Ledbetter

---



Multi-paddock grazing can improve grass quality and production. Credit: Texas A&M AgriLife Research photo

Reducing greenhouse gas emissions in the Southern Great Plains could require a change of grazing management by traditional cow-calf producers, according to a study by Texas A&M AgriLife Research.

Adaptive multi-paddock [grazing](#) is an advanced, more efficient form of rotational grazing and is a potential option to reduce net [greenhouse gas emissions](#), or GHG, on a cow-calf operation in the Southern Great Plains.

This occurs because, compared to continuous grazing, adaptive multi-

paddock grazing results in more carbon sequestration in the soil and produces a higher quality grass that reduces methane production, the study concluded.

The scientific journal, Sustainability, recently published the findings of the study "GHG Mitigation Potential of Different Grazing Strategies in the United States Southern Great Plains."

The research team included Dr. Richard Teague and Dr. Seong Park, both with AgriLife Research, and Stan Bevers, Texas A&M AgriLife Extension Service, all in Vernon, and Dr. Tong Wang, formerly an AgriLife Research post-doctoral researcher now in the South Dakota State department of economics, Brookings, South Dakota.

Ruminants, particularly beef cattle, are perceived by many as a problem since they are a source of greenhouse gas due to the methane produced by rumen fermentation, Park said.

"We believe that conclusion is premature until full ecosystem analyses have outlined the net emissions by considering all emissions compared to carbon sequestration associated with different options in the beef production chain," Teague said.

In this study funded in part by the Dixon Water Foundation, the team considered both GHG emissions and carbon sequestration to calculate net GHG emissions for cow-calf farms grazing only rangeland under different grazing strategies, Park said.

Unlike most published work that isolates the analyses of GHG emission and carbon sequestration, he said they used field-measured soil organic carbon data to estimate the carbon sequestrations for different [grazing management](#) systems.

"Contrary to other publications claiming cow-calf farms are the most significant GHG emission source in the beef production link, our results show that cow-calf farms converting to multi-paddock grazing in the Southern Great Plains region are likely net carbon sinks," Park said. "The continuous grazing was less effective in sequestering carbon."

Beef cattle production management practices in different regions vary greatly in terms of stocking rate, cow size, calving season, forage types and fertilizer use, Bevers said. Analyzing these criteria, the team determined the overall greenhouse gas emissions and main sources of those in the Southern Great Plains region differ from the rest of the U.S. and other countries.

While about one third of U.S. cow-calf only farms are located in the Southern Great Plains, no known net-emission life-cycle assessment study regarding cow-calf production on perennial pastures has been conducted until this one, Wang said.

In this study, the researchers calculated the carbon footprints for cow-calf farms under continuous and multi-paddock grazing strategies using life-cycle assessment modeling, which is a standard assessment of the environmental impacts associated with a wide range of agricultural systems using a "cradle-to-grave" approach.

Compared to the sector approach, which only includes emissions from direct farm activities, life-cycle assessment also includes indirect emissions generated by farm inputs and pre-chain activities, Wang said.

The researchers did find overall GHG emissions are higher in the Southern Great Plains than the other regions, with almost 80 percent of those GHG emissions coming from ruminant digestion.

"But this means there is great potential to reduce these GHG emissions

by increasing grass quality and digestibility using multi-paddock grazing, which could reduce total GHG emissions by as much as 30 percent," Wang said.

Compared to continuous grazing, multi-paddock grazing can improve grass quality as well as grass production, according to ongoing research on this subject.

Teague has studied three grazing management alternatives on neighboring commercial ranches in three proximate counties in north Texas tall grass prairie: continuous grazing with light stocking, representing the best-case scenario for continuous grazing; traditional heavily stocked, continuous grazing, representing the most commonly used grazing management; and adaptive multi-paddock grazing, representing best-case rotational grazing.

"Under multi-paddock grazing management, one paddock is grazed at a time while the other paddocks recover," Teague said. "This grazing strategy uses short periods of grazing, long recovery periods, and adaptively changing recovery periods and other management elements as conditions change."

There is published and anecdotal evidence from producers that, if applied appropriately to produce most advantageous results, multi-paddock grazing can lead to improved water infiltration and soil health, which could lead to increased forage and livestock production, he said.

"Combined with its potential to significantly lower GHG [emissions](#), we conclude that multi-paddock cattle grazing serves as the best carbon mitigation option for this region," Park said.

**More information:** Tong Wang et al. GHG Mitigation Potential of Different Grazing Strategies in the United States Southern Great Plains,

*Sustainability* (2015). [DOI: 10.3390/su71013500](https://doi.org/10.3390/su71013500)

Provided by Texas A&M University

Citation: Cow-calf grazing practices could determine, mitigate greenhouse gas emissions (2015, October 29) retrieved 15 June 2024 from <https://phys.org/news/2015-10-cow-calf-grazing-mitigate-greenhouse-gas.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.