

Dynamic social-network analysis reveals animal social behaviors

October 21 2015



Zebras form large stable groups. Credit: Tanya Berger-Wolf

Communities are defined by flux: friendships that form and break, loyalties that shift, and visitors passing through. But these dynamic interactions aren't represented in static maps of social networks.

Snapshot diagrams—with lines drawn between individuals to show the ties that connect them based on data collected over time—can't tell the whole story.

Zoologists are similarly hampered when trying to analyze the community organization of social animals. The bonding behaviors of related species may seem similar, even though the environments that shaped the animals' group-behavior are quite different.

Two such closely related species are the endangered Grevy's zebra of Africa—the largest surviving wild equid—and the onager, a wild ass of Asia. A new, dynamic social-network analysis tool has revealed that the two species actually have radically different social behaviors and community structures.



The study was published online in *PLOS ONE*.

University of Illinois at Chicago computational ecologist Tanya Berger-Wolf led the multi-disciplinary team that created "CommDy," a dynamic network computational framework, to better understand group behavior and community.

Both Grevy's zebra and the Asian wild asses form what are known as fission/fusion communities, said Berger-Wolf, who is associate professor of computer science at UIC. In fission/fusion communities, individuals meet and spend time with others in different groups at different times.

The two animals' communities look similar, using a traditionally static social-network analysis. But the zebras are few in number, limited in range, and menaced by large predators such as lions. They often lack access to water. Onagers, in contrast, are relatively abundant and widespread, with no major predators and reliable access to water. One

would expect the two species to have evolved very different social structures to cope with their very different circumstances.



A side by side comparison of activity of Grevy's Zebras and Onagers. Credit: Tanya Berger-Wolf (put it all together), Chayant Tanitpathananandh (networks), Daniel Rubenstein (zebras), and Siva Sundaesan (onagers)

To observe the daily interactions within each of the two animal communities, three co-authors on the study from the Mpala Research Centre in Kenya—ecologists Daniel Rubenstein of Princeton University, Siva Sundaesan of the Jackson Hole Conservation Alliance in Wyoming, and Ilya Fischhoff of the University Corporation for Atmospheric Research in Washington, D.C.—drove repeatedly along the same route through the animals' territory and recorded the size, duration and membership of different groups. The new software allowed the researchers to contextualize the observed interactions.

"We're looking for the latent community structure—loyalties, changes in affiliation, visiting with other groups—and the social cost of change," Berger-Wolf said. Some interactions may have a negative impact for an individual, she said, by increasing stress or inviting harassment. Other social contacts may be positive, by increasing status or access to resources.

"The dynamic communities that resulted from that computational analysis were actually strikingly different," Berger-Wolf said. The Grevy's zebra lived in large, stable groups, with loyalty rewarded and visiting with other groups discouraged. Onagers formed smaller, less cohesive groups, with individuals able to change circles with little social cost.

Berger-Wolf said these newly revealed differences make sense, given the

different adaptation each species needed to survive. And for the first time, scientists were able to quantify the differences, using the new computational tool.

Facing a constant threat from a predator, the Grevy's zebra finds strength in numbers, forming large groups of loyal individuals. The large, stable herd is also able to share found resources, like water. The onager, in contrast, can form smaller, more transient groups, being less dependent on the herd for protection or finding scarce resources.

"Dynamic community analysis can be an important tool for testing ideas about the selective ecological and evolutionary forces that generate behaviors, revealing their adaptive value and significance," Berger-Wolf said.

Provided by University of Illinois at Chicago

Citation: Dynamic social-network analysis reveals animal social behaviors (2015, October 21)
retrieved 19 April 2024 from

<https://phys.org/news/2015-10-dynamic-social-network-analysis-reveals-animal.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--