

What are these nanostars in 2Dsuperconductor supposed to mean?





Credit: Moscow Institute of Physics and Technology

Physicists from France and Russia have discovered magnetic disturbances in two-dimensional (2D) layered superconductors resembling small oscillating stars. These star-like electronic excitations are located around individual magnetic atoms within the superconducting material. This experimental observation, made by means of scanning tunneling spectroscopy at only 0.3 degrees above absolute zero, is the direct confirmation of the famous theory by Yu-Shiba-Rusinov, who predicted these quantum-bound magnetic states.



Moreover, it was determined that in the two-dimensional systems, the magnetic excitations spread over longer distances as compared to ordinary three-dimensional superconducting materials. This discovery opens a route for generating more complex quantum states out of chains or clusters of magnetic atoms in superconductors, and which are topologically protected from decoherence. Building and manipulating such protected states is a crucial step toward quantum computers. The results were published in *Nature Physics* this week.

The researchers studied the emergence of Yu-Shiba-Rusinov (YSR) states bound around single magnetic atoms embedded in a twodimensional superconductor. YSR states were theoretically predicted in the 1960s, but little evidence for them has been experimentally revealed until the current study. In the present work, it was found that in twodimensional systems, magnetic excitations extend over a greater distance as compared to ordinary, three-dimensional superconductors, and the emergent YSR quantum states are more stable, which makes them more suitable for a new generation of quantum electronics.

A crystal lattice of a layered superconducting material—niobium diselenide—was used in the tests. With an ultra-low-temperature scanning tunnelling microscope built by Roditchev, the researchers were able to observe for the first time YSR states around single magnetic atoms of iron. "We have demonstrated that the use of two-dimensional superconductors instead of the three-dimensional ones results in an increase in the spatial extension of YSR states for several dozen nanometres, i.e. ten times further than in 'normal,' three-dimensional superconductors. And the area of excitation was shaped like a six-fold electronic 'star' with its rays extending along the axis of the crystal lattice of niobium diselenide. The observed 'stars' are more stable and more suitable to creating new topologically protected states. Non-Abelian anyons can be collected from the YSR state chains, and can then be used as elements in quantum computers of the future," says Vasily Stolyarov,



a co-author of the study and the head of the Laboratory of Topological Quantum Phenomena in Superconducting Systems at MIPT.



Credit: Moscow Institute of Physics and Technology

The experiments described in the article were conducted in Paris. Work is underway at MIPT's Laboratory of Topological Quantum Phenomena in Superconducting Systems to create the experimental conditions necessary to obtain such high-quality results. The laboratory was set up in 2014 using funds from a mega grant awarded to Alexander Golubov, a professor at the University of Twente (Netherlands). The main purpose of the Laboratory is to study the quantum properties of new superconductors and topologically protected materials, and also hybrid artificial systems based on these materials.

The laboratory is being equipped in close cooperation with Roditchev's and Cren's groups and it is expected that the facilities at both laboratories will complement one another.



Yu-Shiba-Rusinov states were predicted in the late 1960s by three physicists from China, the USSR, and Japan independently of one another. <u>They suggested</u> that magnetic atoms introduced into a superconductor must create special states of excitation around themselves—electron-hole standing waves named after their discoverers. Calculations show that areas of topological conductivity may form around these states, in which the current only flows in one direction. Until recently, however, it had not been possible to confirm this prediction experimentally.

For the last 20 years, scientists have been attempting to create quantum systems that will outperform traditional semiconductor-based computers, the development potential of which is now almost exhausted. A number of potential "candidate" systems to be used as a base to build the components of a quantum computer are currently being investigated. The main problem preventing the development of these computers is the high sensitivity of the nanoworld to external influences that destroy quantum states. One promising option is to use topologically protected electron states that are resistant to decoherence. Non-Abelian anyons may be perfect for this; they are not negative ions, but rather special excitations in two-dimensional quantum systems in a magnetic field.

The theory predicts that such non-Abelian anyons may occur in a twodimensional "liquid" of electrons in a superconductor under the influence of a local magnetic field. The electron liquid thus becomes degenerate, i.e. the electrons can have different states at the same energy level. The superposition of several anyons cannot be affected without moving them; therefore, they are completely protected from disturbances.

More information: Gerbold C. Ménard et al. Coherent long-range magnetic bound states in a superconductor, *Nature Physics* (2015). <u>DOI:</u> <u>10.1038/nphys3508</u>



Provided by Moscow Institute of Physics and Technology

Citation: What are these nanostars in 2D-superconductor supposed to mean? (2015, October 16) retrieved 27 June 2024 from <u>https://phys.org/news/2015-10-nanostars-2d-superconductor.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.