

Rising seas will drown mangrove forests

October 14 2015



Mangrove forest at high tide. Credit: Global Change Institute

Mangrove forests around the Indo-Pacific region could be submerged by 2070, international research published today says. Even with relatively low sea-level rises, many mangrove forests had a poor outlook said

Professor Catherine Lovelock, a University of Queensland ecologist.

"Mangrove forests are particularly vulnerable," she said.

"Mangroves are predicted to be submerged in parts of Thailand, Sumatra, Java, Papua New Guinea and the Solomon Islands."

However the outlook in other parts of the world was more positive.

"Our modelling shows mangroves are likely to persist in east Africa, the Bay of Bengal, eastern Borneo and north-western Australia - areas where there are relatively large tidal ranges and/or higher sediment supply," said Professor Lovelock, who works in UQ's School of Biological Sciences and the Global Change Institute.

"Even in other areas though, the good news is that through accretion of sediment and maintenance of wetland soils, [mangrove forests](#) do have the capacity to avoid inundation and keep pace with [sea-level rise](#)."

Professor Lovelock said this was why there was an urgent need to plan for the maintenance of sediment supply in river systems that were likely to be dammed or heavily modified in future.

"The Indo-Pacific region holds most of the world's mangrove forests, but sediment delivery in this region is declining, due to activities such as dam construction," she said.

"This is of particular concern as this region is expected to have variable but high rates of future sea-level rise."

Forest degradation had to be reversed because it reduced organic inputs to soils that was vital for mangrove survival, Professor Lovelock said.

Plans should be made for the landward migration of vulnerable mangrove forests to higher elevations.

"Intertidal mangrove forests occur on tropical and subtropical shorelines, and provide a wide range of ecosystem services - to fisheries, in coastal protection and in carbon sequestration - with an estimated value of \$USD194,000 per hectare per year," she said.

"Sea-level rise could threaten the long-term sustainability of coastal communities and of valuable ecosystems such as coral reefs, salt marshes and mangroves.

Professor Lovelock said the researchers analysed trends based on data from an international network of 27 sites.

UQ's Global Change Institute and the Australian Research Council funded the project.

The vulnerability of Indo-Pacific mangrove forests to sea-level rise is published today in *Nature*.

More information: *Nature*, [DOI: 10.1038/nature15538](https://doi.org/10.1038/nature15538)

Provided by University of Queensland

Citation: Rising seas will drown mangrove forests (2015, October 14) retrieved 26 April 2024 from <https://phys.org/news/2015-10-seas-mangrove-forests.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.