

Speedy evolution affects more than one species

October 22 2015



A parasitic wasp (*Diachasma alloeum*) that preys upon a fruit fly species known as an apple maggot. Researchers found that the wasp evolved after the fruit fly evolved when it changed its mating and egg-laying habits. Photo courtesy of James Smith, Lyman Briggs College.

The concept that biodiversity feeds upon itself is not uncommon in the

world of evolution. The problem is a lack of hard data that shows this process to be naturally occurring.

However, recent research by a team of scientists, including a Michigan State University entomologist, finds that recent evolutionary changes - in this case in a [new species](#) of fruit fly - have an almost domino effect on a number of species.

The research, published this week in the *Proceedings of the National Academy of Sciences*, follows up work done by the team several years ago that found changes in mating habits resulted not only in a new species of fruit fly, but also led to a new species of the parasitic wasps that prey on them.

"The new study extends the earlier work by showing that new fruit fly species provide suitable habitat not just for one new parasitoid species, but for multiple new species," said James Smith, an MSU entomologist and professor in Lyman Briggs College.

The [fruit flies](#) in question evolved into new species when they began laying their eggs and mating on apple trees, as opposed to their native hawthorn tree hosts. Three different kinds of [parasitoid wasps](#) were collected from a number of different fly [host plant](#) environments in the wild.

Analyses in the lab showed that all three of the different kinds of wasps had diverged from others of the same kind, both genetically and with respect to host-associated physiology and behavior.

"In a sense," Smith said, "they have caught an entire community of parasitoids actively ecologically diverging in response to a historically documented host plant shift of their fly host."

These evolutionary changes, known as "sequential" or "cascading" events, may provide additional information helping explain why some groups of organisms, such as plants, the insects that feed on them and the parasites that attack the insects, are more diverse and species-rich than other groups.

"Why are there so many insect species?" Smith asked. "Speciation cascades provide one explanation for how a lot of [species](#) might be generated in a relatively short period of time."

More information: Sequential divergence and the multiplicative origin of community diversity,
www.pnas.org/cgi/doi/10.1073/pnas.1424717112

Provided by Michigan State University

Citation: Speedy evolution affects more than one species (2015, October 22) retrieved 20 March 2024 from <https://phys.org/news/2015-10-speedy-evolution-affects-species.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--