


Unraveling the complex, intertwined electron phases in a superconductor

October 26 2015



A rendering of the charge density wave. This pattern of dumbbells, each representing the electronic orbital on an oxygen atom, may look static, but if you look closely you'll see that the colors of the orbitals change as you move from left to right. This color change (yellow, white, blue, white, yellow, white, blue...) represents changes in charge density (high, medium, low, medium, high, medium, low...) moving across the plane from left to right. Credit: Brookhaven

A team led by researchers from the U.S. Department of Energy's Brookhaven National Laboratory and Cornell University has characterized a key arrangement of electrons in a high-temperature superconductor, a material that can conduct electricity with almost no energy loss without being ultra-chilled. The material is a member of a family of copper-oxygen-based superconducting compounds—the cuprates—that are prime candidates for numerous potential high-impact applications, including extremely efficient electricity generation, storage, and transmission across the nation's power grid.

The phenomenon they studied is known as an electron density wave. Unlike the other electrons in the material, which move about freely, the density wave is a periodic, fixed electron phase that seems to compete with and hinder the superconducting phase. Many researchers believe that the density wave is the key to unlocking the cuprates: If they can thoroughly understand the electron density wave, they say, they may be able to determine how to suppress or remove it to induce superconductivity, possibly even at room temperature. But to achieve this goal, they first must gain a thorough understanding of the causes of the electron density wave.

This study, published in the October 26, 2015, online edition of *Nature Physics*, is the first to identify the atomic-scale origins and influences that produce the density wave in cuprates.

"In a metal, you have electrons going in every direction, with a wide range of energies and momenta. It's almost like a gas of electrons," said the study's lead scientist, Séamus Davis, a physicist at Brookhaven and the J.G. White Distinguished Professor of Physical Sciences at Cornell.

"Which of these electrons provide the components that make up the density wave state in cuprates? We were able to answer this question. We identified both the energy and momenta of the relevant electrons."

Another piece of the cuprate puzzle

There are several known members of the cuprate family. The "best" one discovered so far works as a superconductor at about 140 Kelvin (-130 Celsius), roughly halfway between absolute zero (-273°C) and room temperature (21°C). All have layered structures, with superconductivity taking place in the crystalline copper-oxide (CuO₂) layers, each just one molecule thick. Aside from the superconducting phase, these materials exhibit another mysterious form of electronic order, dubbed the "pseudogap" phase. While very poorly understood in general, its characteristic energy scale, called an energy gap, is well established. Davis' group and others have studied this phase extensively.

Last year, along with researchers led by Brookhaven/Columbia University School of Engineering physicist Simon Billinge, the team established the first firm link between the disappearance of the density wave within the [pseudogap phase](http://phys.org/news/2014-05-solution-long-standing-mysteries-cuprate-superconductivity.html) and the emergence, as stated by Davis, of "universally free-flowing electrons needed for unrestricted superconductivity" [see: <http://phys.org/news/2014-05-solution-long-standing-mysteries-cuprate-superconductivity.html>].

In this most recent work, led at Brookhaven by physicist Kazuhiro Fujita, the most important result seems to suggest another intimate link between the electron density wave and the pseudogap phase: The characteristic energy of the density wave's modulation is equal to the pseudogap energy. Moreover, they showed that the electrons making up the electron density wave have the characteristic momentum at which the pseudogap appears. The fundamental implication is that the

pseudogap state is somehow the "parent" of the density wave state.

Imaging electrons with electrons

The novel experimental approach and the concept of the study itself were conceived after one of the paper's authors, Harvard University theoretical physicist Subir Sachdev, pondered the differing electron density wave behavior along the different bonds of the CuO₂ unit cell.

"In looking at some of the past data taken by Davis's group, I noticed that the electron behavior was unique to the horizontal bonds of the CuO₂ plane," he said. "I wondered if this could be the charge density wave."

He contacted Davis and they began designing the experiment. They proposed a new way to study a cuprate, one that no other group had tried: a powerful imaging technique developed by Davis, called sublattice imaging-which is performed using a specialized scanning tunneling microscope (STM) capable of determining the electronic structure in different subsets of the atoms in the crystal, the so-called sublattices.

An STM images a sample using electrons; the type used here can "see" details smaller than the unit cell of the CuO₂ crystal. It uses a very sharp metallic tip, placed extremely close to the sample surface-less than a tenth of a nanometer away. A small voltage is then applied across the tip and sample, which, because of their proximity, allows electrons to "tunnel" through the vacuum gap between the tip and sample. The rate of electrons passing through the tip-in other words, the current-is a measure of the density of [electrons](#) in the tiny region of the sample directly below it. So as the tip is passed across the sample surface, it creates a spatial image of the [electron density](#) and, separately, of the crystal lattice.

More information: Atomic-scale electronic structure of the cuprate d-symmetry form factor density wave state, *Nature Physics*, [DOI: 10.1038/nphys3519](https://doi.org/10.1038/nphys3519)

Provided by Brookhaven National Laboratory

Citation: Unraveling the complex, intertwined electron phases in a superconductor (2015, October 26) retrieved 18 April 2024 from <https://phys.org/news/2015-10-unraveling-complex-intertwined-electron-phases.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.