

Bird decline shows that climate change is more than just hot air

November 16 2015

Scientists have long known that birds are feeling the heat due to climate change. However, a new study of a dozen affected species in the Western Cape suggests their decline is more complex than previously thought – and in some cases more serious.

According to the study, published today (Monday) in Conservation Physiology, by scientists from the Percy Fitzpatrick Institute at the University of Cape Town and Nelson Mandela Metropolitan University, there could be several reasons why <u>birds</u> are being negatively affected by man-made climate change.

They suggest that contrary to expectations the birds' heat tolerance – or lack thereof—is not necessarily the main factor chasing <u>species</u> out of their preferred habitat. Other factors like changing fire and rainfall patterns, and new bird behaviour patterns could also be responsible for the decline, according to the study which includes some well-known species such as the Malachite Sunbird and the Familiar Chat.

Lead researcher Susie Cunningham says a better understanding of how climate change affects <u>bird species</u> could help develop conservation strategies to increase chances of survival:

"We know climate change is linked to changes in species' numbers and distributions, but we don't always know exactly how or why," Cunningham says. "We need to figure out the factors actually driving declines before we can develop proper conservation measures to halt



them."

"Fynbos birds are particularly important in this regard because they live in an area that has been extremely stable, climatologically speaking, for a very long time. So changes in climate are not something they are used to. Furthermore, six of the species we studied are endemic to the fynbos, so if we lose them from this biome, we lose them altogether," she explains.

The new study profiles the well-being of twelve fynbos bird species based on a comparison of data contained in two Southern Africa Bird Atlas surveys conducted fifteen years apart. This data was matched with climate data for the comparative period, as well as with physiological data. A key aspect of the study is a comparison of climate and bird population data with the heat response – or 'thermal tolerance' – of each bird species. In this way researchers assessed to what extent birds can cope with changing temperature, and whether this is the primary indicator of changes in bird abundance.

To obtain bird physiological data researchers subjected birds to a range of temperature tests to determine their physiological response. Considerable variation in the relationship between <u>physiological</u> <u>response</u> to temperature and rate of species' decline prevented researchers confirming a direct link across the board between how species are faring under current climate change, and their thermal physiology.

Despite this, the study shows 'striking similarity' between bird declines and increases in average temperature, most notably in two bird species occupying cooler rangelands – the Cape Rockjumper and the Protea Seed-eater. Reporting rates of these two species dropped by 31 percent and 32 percent respectively in the study area during the period under review.



In addition the Cape Rockjumper is also the only species whose decline is clearly linked to a low physiological tolerance of heat, prompting the study authors to call for urgent conservation intervention: "Conservation action appears to be urgently needed for the Cape Rockjumper and Protea Seed-eater if their declines are to be properly understood and halted," the study says. "We need further research to determine the cause of decline of Protea Seed-eaters, and to assess how Cape Rockjumpers may best be assisted to cope in the face of continuing climate change."

Cunningham says the overall findings suggest scientists should not jump to conclusions about bird physiology when assessing the effects of <u>climate change</u>: "The main findings are that physiology, though often considered the ultimate factor limiting species distributions, may not be the factor responsible for warming-related declines in most Fynbos birds," she explains.

More information: "The role of thermal physiology in recent declines of birds in a biodiversity hotspot" Robyn Milne, Susan J. Cunningham, Alan T. K. Lee, and Ben Smit, *Conservation Physiology*, <u>DOI:</u> <u>10.1093/conphys/cov048</u>

Provided by Oxford University Press

Citation: Bird decline shows that climate change is more than just hot air (2015, November 16) retrieved 6 May 2024 from <u>https://phys.org/news/2015-11-bird-decline-climate-hot-air.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.