

# Early galaxies more efficient at making stars, Hubble survey reveals

November 19 2015

---



This image shows a region of the CANDELS GOODS-South field, which is one of the fields used in this study. This image combines data taken from Hubble Space Telescope's optical and near-infrared cameras, and contains galaxies at a range of distances. The larger galaxies are relatively close by, while the smallest specks hail from the earlier universe. Some of the smallest dots in this image are those used in this study; their light is coming from 0.5 to 1.5 billion years after the Big Bang. Credit: NASA, ESA, A. Koekemoer and the CANDELS science team

A study published in today's *Astrophysical Journal* by University of Texas at Austin assistant professor Steven Finkelstein and colleagues reveals that galaxies were more efficient at making stars when the universe was younger. The announcement explains the team's discovery, announced in the journal's September 1 issue, that there are a lot more bright, highly star-forming galaxies in the early universe than scientists previously thought.

"This was an unexpected result," Finkelstein said. "It has implications for galaxy formation at the earliest times" in the universe.

For both studies, his team used galaxy observations from Hubble Space Telescope's CANDELS survey, of which he is a team member. Hubble's largest survey to date, CANDELS stands for Cosmic Assembly Near-infrared Deep Extragalactic Legacy Survey.

Today's finding stems from studies of about 8,000 CANDELS galaxies seen at times ranging from 0.75 to 1.5 billion years after the Big Bang (that is, between redshift four and redshift seven). As the universe is a little less than 14 billion years old, this corresponds to only the first five to 10 percent of the history of the universe.

The team deduced the rate of [star formation](#) in these galaxies from the Hubble images, by noting their brightness in ultraviolet light, and then correcting this measurement depending on how much light-absorbing dust the galaxy contains. The dust estimation comes from the Hubble images, too. The redder a galaxy is, the dustier it is.

Investigating the highly star-forming galaxies further, they compared the mass in stars in these galaxies to the theoretically predicted rate at which galaxies grow their mass in the early universe. They found higher masses than predicted, implying that galaxies are more efficient at turning gas into stars in the early universe than they are today.

There could be a couple of different reasons why, Finkelstein said.

First, as the universe has been expanding outward since the Big Bang, at earlier times everything in the universe was packed closer together, including the gas in galaxies. Dense gas is the material that makes stars, so perhaps these galaxies simply had more of it.

Second: feedback. "No galaxy is 100 percent efficient at turning gas into stars," Finkelstein said, explaining that there are several mechanisms inside galaxies that can cause some of the gas to not form stars. These include things like the massive explosions called supernovae, winds from massive stars, and active supermassive black holes that can heat their surrounding gas. Altogether, these barriers to star formation collectively are called "feedback." Finkelstein said that galaxies at earlier times may experience less feedback, and so may form [stars](#) more readily.

He anticipates that these bright galaxies in the [early universe](#) can be studied in greater detail with the forthcoming James Webb Space Telescope (JWST), the infrared successor to Hubble, which will launch in 2018. Future studies with JWST should provide a better understanding of star formation in early [galaxies](#).

**More information:** \* "An Increasing Stellar Baryon Fraction in Bright Galaxies at High Redshift," Steven L. Finkelstein et al., 2015 December 1, *Astrophysical Journal* [iopscience.iop.org/article/10.1088/0004-637X/814/2/95](http://iopscience.iop.org/article/10.1088/0004-637X/814/2/95) , Arxiv: [arxiv.org/abs/1504.00005](http://arxiv.org/abs/1504.00005)

\* "The Evolution of the Galaxy Rest-frame Ultraviolet Luminosity Function Over the First Two Billion Years," Steven L. Finkelstein et al., 2015 September 1, *Astrophysical Journal* [iopscience.iop.org/article/10.1088/0004-637X/810/1/71](http://iopscience.iop.org/article/10.1088/0004-637X/810/1/71) , Arxiv: [arxiv.org/abs/1410.5439](http://arxiv.org/abs/1410.5439)

Provided by University of Texas at Austin

Citation: Early galaxies more efficient at making stars, Hubble survey reveals (2015, November 19) retrieved 13 March 2024 from <https://phys.org/news/2015-11-early-galaxies-efficient-stars-hubble.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.