

Great Barrier Reef protecting against landslides, tsunamis

November 25 2015



Landslide distribution and geomorphology. Credit: Webster et al, University of Sydney

The world-famous Australian reef is providing an effective barrier against landslide-induced tsunamis, new research shows.

What has developed into the Great Barrier Reef was not always a barrier



reef - it was once a fringing reef and did not offer the same protective quality. This is because the coast at this time was much closer to the source of the tsunamis, said lead author of the paper, Associate Professor Jody Webster, from the Geocoastal Research Group at the University of Sydney.

The research shows a shallow underwater landslide occurred 20,000-14,000 years ago, which caused a tsunami 2-3m high. The tsunami could have impacted Aborigines living at the time along estuaries and on islands off the paleo-coastline, which has since receded under the rising sea levels that followed the last ice age.

The 7km-wide landslide occurred off the edge of the continental shelf causing the tsunami on the paleo-coastline lying between Airlie Beach and Townsville in the northern State of Queensland.

Details of the discovery of the submarine landslide and tsunami were published this week in *Marine Geology*. The international team of researchers used sophisticated computer simulations to recreate what the tsunami would have looked like.

Associate Professor Webster said similar landslides under the sea could occur without our knowledge.

"There is a relatively low chance that a similar submarine landslide with the potential to cause a tsunami of up to three metres or more would happen today," Associate Professor Webster said.



Fig. 6. Landslide and tsunami simulations



Landslide and tsunami simulations at 0m, -50m and -70m sea-level scenarios. Credit: Webster et al, University of Sydney

"However, if one did occur, our findings suggest that the Great Barrier Reef is doing us a great service because of its ability to absorb some of that potential wave energy."

Just how much energy would be absorbed and what the extent of damage could be done by rising sea levels and tsunamis or king tides is the subject of future research.

In reaching their findings, Dr Jon Hill from the University of York created visual simulations of the <u>tsunami</u> impact at today's <u>sea level</u>, as well as at a depth of -70m, where the paleo-coastline was before it



receded to its current position and was replaced at the shelf edge by the formation of the Great Barrier Reef.

The research team has named the submarine landslide the Viper Slide because of its location adjacent to Viper Reef.

"The discovery of the Viper Slide is the first solid evidence that submarine landslides existed on the Great Barrier Reef," said Dr Robin Beaman from James Cook University - a member of the expedition that mapped the slide.

More information: Jody M. Webster et al. Submarine landslides on the great barrier reef shelf edge and upper slope: A mechanism for generating tsunamis on the north-east Australian coast?, *Marine Geology* (2015). DOI: 10.1016/j.margeo.2015.11.008

Provided by University of Sydney

Citation: Great Barrier Reef protecting against landslides, tsunamis (2015, November 25) retrieved 3 May 2024 from <u>https://phys.org/news/2015-11-great-barrier-reef-landslides-tsunamis.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.