

# Neonicotinoid pesticides linked to butterfly declines in the UK

November 24 2015, by Jacqui Bealing

---



The use of neonicotinoid pesticides may be contributing to the decline of butterflies in the UK, according to a new study involving the University of Sussex.

Previous studies have demonstrated that these chemicals appear to be harming bees, birds and other wildlife.

The new study is the first [scientific evidence](#) of a possible negative impact on widespread UK [butterflies](#).

The chemicals remain in the environment and can be absorbed by the wildflowers growing in field margins, many of which provide a nectar source for butterflies and food-plants for their caterpillars.

Researchers at the Universities of Stirling and Sussex, in partnership with Butterfly Conservation and the Centre for Ecology and Hydrology, found population trends of 15 species showed declines associated with neonicotinoid use, including Small Tortoiseshell, Small Skipper and Wall species.

The study, published today (24 November 2015) in the journal *PeerJ*, is based on data gathered by volunteers from more than 1000 sites across the UK as part of the long-running UK Butterfly Monitoring Scheme (UKBMS).

Dave Goulson, Professor of Biology at the University of Sussex said: "Many of us can remember a time when our meadows and hedgerows had far more butterflies, bees and other insects than today. This study adds to the growing mountain of evidence that neonicotinoids are one of the causes of these declines."

Ecologist Dr Andre Gilburn of the University of Stirling, who led the study, said: "Our study not only identifies a worrying link between the use of neonicotinoids and declines in butterflies, but also suggests that the strength of their impact on many species could be huge."

Dr Tom Brereton, Head of Monitoring at Butterfly Conservation, said: "We are extremely concerned with the findings of the study and are calling for urgent research to see whether the correlations we found are caused by neonicotinoid use, or some other aspect of intensive farming. Widespread butterflies have declined by 58 percent on farmland in England over the last 10 years, giving concern for the general health of the countryside and for these and other insects in particular."

**More information:** Andre S. Gilburn et al. Are neonicotinoid insecticides driving declines of widespread butterflies?, *PeerJ* (2015).  
[DOI: 10.7717/peerj.1402](https://doi.org/10.7717/peerj.1402)

Provided by University of Sussex

Citation: Neonicotinoid pesticides linked to butterfly declines in the UK (2015, November 24)  
retrieved 10 April 2024 from  
<https://phys.org/news/2015-11-neonicotinoid-pesticides-linked-butterfly-declines.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--