


Increasing production of seed oils

November 9 2015



WT: wild-type seed. FUS3pro::WRI1: seed of the transgenic plant with extended oil synthesis phase during seed formation. FUS3pro::WRI1/12s1.4: seed of the transgenic plant with extended oil synthesis phase and disrupted formation of major seed storage proteins. Credit: NIBB

Plant-derived oils are widely used all over the world both for food and for industrial purposes. In recent years they have also attracted attention as raw materials for potential bio-fuels and bio-plastics that are friendly to the environment. Because of these, the demand for vegetable oils is expanding year by year. Researchers in Japan have succeeded in inducing the genes involved in oil synthesis in seeds to work for longer



periods of time, thereby allowing them to accumulate more seed oil. Their results have been reported online in the *Plant Biotechnology Journal* on October 26, 2015.

Many plants store oil and protein in their <u>seeds</u> as energy sources needed for germination. The National Institute for Basic Biology's Drs. Masatake Kanai, Shoji Mano and Mikio Nishimura et al., in the process of studying the mechanism that controls the synthesis and degradation of oils in <u>plant seeds</u>, revealed that oils are actively synthesized only during the mid-phase of the seed formation process.

The research group considered that by extending the period seeds synthesize oils, they should be able to increase their oil content in seeds. Therefore, using the model plant *Arabidopsis*, they extended the expression time of WRI1, a gene that activates oil synthesis. As a result, they were able to increase the oil content of the seeds to 140% as compared to wild-type. Also, while extending the oil synthesis phase, suppressing protein synthesis in the late-phase of seed formation increased the oil content of the seeds as compared to wild-type to 170%.

Because of this study it is now clear that the length of the oil synthesis phase in seed formation is one of the primary factors that determine final oil content. Moreover, by suppressing protein synthesis while extending the oil synthesis period they succeeded in a further large increase in oil production.

Dr. Kanai of the research group, said "By applying the current results to crop breeding, prolonging the period of oil synthesis will generate many new crops with high <u>oil content</u>".

More information: Masatake Kanai et al. Extension of oil biosynthesis during the mid-phase of seed development enhances oil content in seeds, *Plant Biotechnology Journal* (2015). DOI:



10.1111/pbi.12489

Provided by National Institutes of Natural Sciences

Citation: Increasing production of seed oils (2015, November 9) retrieved 19 April 2024 from https://phys.org/news/2015-11-production-seed-oils.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.