

Sharks' hunting ability destroyed under climate change

November 12 2015



Great white shark at Isla Guadalupe, Mexico, August 2006. Credit: Terry Goss/Wikipedia

The hunting ability and growth of sharks will be dramatically impacted by increased CO₂ levels and warmer oceans expected by the end of the century, a University of Adelaide study has found.

Published today in the journal *Scientific Reports*, marine ecologists from

the University of Adelaide's Environment Institute report long-term experiments that show warmer waters and [ocean acidification](#) will have major detrimental effects on sharks' ability to meet their energy demands, with the effects likely to cascade through entire ecosystems.

The laboratory experiments, studying Port Jackson sharks and including large tanks with natural habitat and prey, found embryonic development was faster under elevated temperatures. But the combination of warmer water and high CO₂ increased the sharks' energy requirement, reduced metabolic efficiency and removed their ability to locate [food](#) through olfaction (smelling). These effects led to marked reductions in growth rates of sharks.

"In warmer water, sharks are hungrier but with increased CO₂ they won't be able to find their food," says study leader Associate Professor Ivan Nagelkerken, Australian Research Council (ARC) Future Fellow.

"With a reduced ability to hunt, sharks will no longer be able to exert the same top-down control over the [marine food webs](#), which is essential for maintaining healthy ocean ecosystems."

PhD student Jennifer Pistevos, who carried out the study, says the Port Jackson is a bottom-feeding shark that primarily relies on its ability to smell to find food. Under higher CO₂, the sharks took a much longer time to find their food, or didn't even bother trying, resulting in considerably smaller sharks.

Most research studying the effects of ocean acidification and climate change on fish behaviour has concentrated on small fish prey. Long-term studies on the behaviour and physiology of large, long-lived predators are largely lacking.

Fellow University of Adelaide marine ecologist Professor Sean Connell

says the results of the study provide strong support for the call to prevent global overfishing of [sharks](#).

"One-third of shark and ray species are already threatened worldwide because of overfishing," Professor Connell says. "Climate change and ocean acidification are going to add another layer of stress and accelerate those extinction rates."

Provided by University of Adelaide

Citation: Sharks' hunting ability destroyed under climate change (2015, November 12) retrieved 23 April 2024 from <https://phys.org/news/2015-11-sharks-ability-climate.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.