

Thermal sensitivity of marine communities reveals the most vulnerable to global warming

November 11 2015



Reef Life Survey site in Rana, Indonesia. Credit: Rick Stuart-Smith

The sensitivity of marine communities to ocean warming rather than

rising ocean temperatures will have strong short-term impacts on biodiversity changes associated with global warming, according to new research.

The study, involving scientists from the Universities of Southampton (UK), Stockholm (Sweden) and Tasmania (Australia), found that while some [communities](#) - a set of interacting [species](#) - are comprised of species that live at the edge of how much heat they can tolerate, others may have a greater tolerance or 'bias' to warming oceans. This bias is due to the combination of historical, ecological and evolutionary factors that play an important role in determining which species are found where.

By understanding the size of a communities' thermal bias, scientists will be able to predict and prioritise which areas are going to be most vulnerable to ocean warming.

Study co-author Dr Amanda Bates, from Ocean and Earth Science at the University of Southampton, said: "In 100 years from now, 100 per cent of species in many communities will be lost and replaced by new species able to tolerate warmer conditions, leading to a redistribution of species across the globe."

Using data from the [Reef Life Survey](#), the researchers measured the geographical and thermal distributions of 2,695 shallow reef fish and 1,225 invertebrates from Greenland to Australia to measure the thermal bias for communities around the world.

They found that locations where the average summer [sea surface temperature](#) is presently 24 °C, such as the Gulf of Thailand, southwestern Caribbean and Three Kings-North Cape in New Zealand, are the most vulnerable to changing community biodiversity. This is because most of the species making up these communities are already living near the edge of their temperature distribution.

Dr Bates said: "A strong focus in climate change ecology has been on quantifying the exposure of different regions of the globe to warming. Our work offers new tools for measuring the sensitivity of communities to change including accurate indicators that can be used to predict vulnerability."

The research, which is published in the journal *Nature*, was funded by the Australian Research Council and Marine Biodiversity Hub.

More information: Rick D. Stuart-Smith et al. Thermal biases and vulnerability to warming in the world's marine fauna, *Nature* (2015).

[DOI: 10.1038/nature16144](https://doi.org/10.1038/nature16144)

Provided by University of Southampton

Citation: Thermal sensitivity of marine communities reveals the most vulnerable to global warming (2015, November 11) retrieved 2 May 2024 from

<https://phys.org/news/2015-11-thermal-sensitivity-marine-reveals-vulnerable.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--