

Scientists discover that a single layer of tiny diamonds increases electron emission 13,000-fold

December 10 2015



Nick Melosh, an associate professor at SLAC and Stanford, holds a model of a diamondoid. Credit: SLAC National Accelerator Laboratory

They sound like futuristic weapons, but electron guns are actually



workhorse tools for research and industry: They emit streams of electrons for electron microscopes, semiconductor patterning equipment and particle accelerators, to name a few important uses.

Now scientists at Stanford University and the Department of Energy's SLAC National Accelerator Laboratory have figured out how to increase these electron flows 13,000-fold by applying a single layer of diamondoids – tiny, perfect diamond cages – to an electron gun's sharp gold tip.

The results, published today in *Nature Nanotechnology*, suggest a whole new approach for increasing the power of these devices. They also provide an avenue for designing other types of electron emitters with atom-by-atom precision, said Nick Melosh, an associate professor at SLAC and Stanford who led the study.

Diamondoids are interlocking cages made of carbon and hydrogen atoms. They're the smallest possible bits of diamond, each weighing less than a billionth of a billionth of a carat. That small size, along with their rigid, sturdy structure and high chemical purity, give them useful properties that larger diamonds lack.

SLAC and Stanford have become one of the world's leading centers for diamondoid research. Studies are carried out through SIMES, the Stanford Institute for Materials and Energy Sciences, and a lab at SLAC is devoted to extracting diamondoids from petroleum.

In 2007, a team led by many of the same SIMES researchers showed that a single layer of diamondoids on a metal surface could emit and focus <u>electrons</u> into a tiny beam with a very narrow range of energies.





The research team used tiny nanopillars of germanium wire as stand-ins for the tips of electron guns in experiments aimed at improving electron emission. This image was made with a scanning electron microscope – one of a number of devices that use emitted electrons. Credit: Karthik Narasimha/Stanford

The new study looked at whether a diamondoid coating could also improve emissions from electron guns.



One way to increase the power of an electron gun is to make the tip really sharp, which makes it easier to get the electrons out, Melosh said. But these sharp tips are unstable; even tiny irregularities can affect their performance. Researchers have tried to get around this by coating the tips with chemicals that boost <u>electron emission</u>, but this can be problematic because some of the most effective ones burst into flames when exposed to air.

For this study, the scientists used tiny nanopillars of germanium wire as stand-ins for electron gun tips. They coated the wires with gold and then with diamondoids of various sizes.





Germanium nanopillars were coated with gold and then with diamondoids of various sizes. The scientists got the best results by coating the pillars with diamondoid molecules that consist of four "cages;" this increased the emission of electrons from the tips 13,000-fold. Credit: Karthik Narasimha/Stanford

When the scientists applied a voltage to the nanowires to stimulate the release of electrons from the tips, they found they got the best results from tips coated with diamondoids that consist of four "cages." These released a whopping 13,000 times more electrons than bare gold tips.

Further tests and computer simulations suggest that the increase was not due to changes in the shape of the tip or in the underlying gold surface. Instead, it looks like some of the diamondoid molecules in the tip lost a single electron – it's not clear exactly how. This created a positive charge that attracted electrons from the underlying surface and made it easier for them to flow out of the tip, Melosh said.

"Most other molecules would not be stable if you removed an electron; they'd fall apart," he said. "But the cage-like nature of the diamondoid makes it unusually stable, and that's why this process works. Now that we understand what's going on, we may be able to use that knowledge to engineer other materials that are really good at emitting electrons."





Diamondoid structures tested in the experiment; the two on the bottom, which consist of four "cages" with carbon atoms at each corner, produced the biggest gains in electron emission. The chemical tags at the bottom of each molecule were added to help the diamondoids stick to the gold surface of the nanopillars. Credit: Karthik Narasimha/Stanford





SIMES researchers Nick Melosh, left, and Jeremy Dahl in a Stanford laboratory with equipment used to perform diamondoid experiments. Credit: SLAC National Accelerator Laboratory

More information: Karthik Thimmavajjula Narasimha et al. Ultralow effective work function surfaces using diamondoid monolayers, *Nature Nanotechnology* (2015). DOI: 10.1038/nnano.2015.277

Provided by SLAC National Accelerator Laboratory



Citation: Scientists discover that a single layer of tiny diamonds increases electron emission 13,000-fold (2015, December 10) retrieved 1 May 2024 from https://phys.org/news/2015-12-scientists-layer-tiny-diamonds-electron.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.