

Astronomers find new way to measure the pull of gravity at the surface of distant stars

January 1 2016



How much would you weigh on another star? The timescale of turbulence and



vibration at a star's surface, based on its brightness variations, tells you its surface gravity. If stars had solid surfaces on which you could stand, then your weight would change from star to star. Here we show how much a 75-kg adult would tip the bathroom scale in the surface gravities of three stars. The sun is hotter than a sauna, but don't expect to lose weight there. You'd weigh 20 times more than on Earth. A red giant star (the far-future fate of our Sun, with a diameter about 35 times larger) has a much weaker pull at its surface, so you'd be 50 times lighter. Credit: Jaymie Matthews and Thomas Kallinger

Researchers have found a new way to measure the pull of gravity at the surface of a star. For distant stars with planets orbiting them, this information is key in determining whether any of those planets can harbour life.

The new method is described in a study published today in *Science Advances*. The research was led by University of Vienna's Thomas Kallinger and involved UBC Professor Jaymie Matthews as well as astronomers from Germany, France and Australia.

Knowing the <u>surface gravity</u> of a star is essentially knowing how much you would weigh on that star. If stars had solid surfaces on which you could stand, then your weight would change from star to star. The Sun is hotter than a sauna, but don't expect to lose weight there. You'd weigh 20 times more than on Earth. A red giant star (the far-future fate of our Sun) has a much weaker pull at its surface, so you'd be 50 times lighter.

The new method allows scientists to measure surface gravity with an accuracy of about four per cent, for stars too distant and too faint to apply current techniques. Since surface gravity depends on the star's mass and radius (just as your weight on Earth depends on its mass and radius), this technique will enable astronomers to better gauge the masses and sizes of distant stars. It will play an exciting role in the study of



<u>planets</u> beyond the Solar System, many so distant that even the basic properties of the stars they orbit can't be measured accurately.

"If you don't know the star, you don't know the planet," said study coauthor, UBC Professor Jaymie Matthews. "The size of an exoplanet is measured relative to the size of its parent star. If you find a planet around a star that you think is Sun-like but is actually a giant, you may have fooled yourself into thinking you've found a habitable Earth-sized world. Our technique can tell you how big and bright is the star, and if a planet around it is the right size and temperature to have water oceans, and maybe life."

The new technique called the autocorrelation function timescale technique, or timescale technique for short, uses subtle variations in the brightness of <u>distant stars</u> recorded by satellites like Canada's MOST and NASA's Kepler missions.

Future space satellites will hunt for planets in the 'Goldilocks Zones' of their stars. Not too hot, not too cold, but just right for liquid water oceans and maybe life. Future exoplanet surveys will need the best possible information about the stars they search, if they're to correctly characterize any planets they find.

"The timescale technique is a simple but powerful tool that can be applied to the data from these searches to help understand the nature of stars like our Sun and to help find other planets like our Earth," said Kallinger, the study's lead author.

More information: Precise stellar surface gravities from the time scales of convectively driven brightness variations, <u>dx.doi.org/10.1126/sciadv.1500654</u>



Provided by University of British Columbia

Citation: Astronomers find new way to measure the pull of gravity at the surface of distant stars (2016, January 1) retrieved 1 May 2024 from <u>https://phys.org/news/2016-01-astronomers-gravity-surface-distant-stars.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.