

## Dark 'noodles' may lurk in the Milky Way

January 21 2016



CSIRO's Compact Array in Australia is shown under the night lights of the Milky Way. Credit: Alex Cherney

Invisible structures shaped like noodles, lasagne sheets or hazelnuts could be floating around in our Galaxy radically challenging our understanding of gas conditions in the Milky Way.

CSIRO astronomer and first author of a paper released in *Science* Dr Keith Bannister said the structures appear to be 'lumps' in the thin gas that lies between the stars in our Galaxy.



"They could radically change ideas about this interstellar gas, which is the Galaxy's star recycling depot, housing material from old stars that will be refashioned into new ones," Dr Bannister said.

Dr Bannister and his colleagues described breakthrough observations of one of these 'lumps' that have allowed them to make the first estimate of its shape.

The observations were made possible by an innovative new technique the scientists employed using CSIRO's Compact Array telescope in eastern Australia.

Astronomers got the first hints of the mysterious objects 30 years ago when they saw radio waves from a bright, <u>distant galaxy</u> called a quasar varying wildly in strength.

They figured out this behaviour was the work of our Galaxy's invisible 'atmosphere', a thin gas of electrically charged particles which fills the space between the stars.

"Lumps in this gas work like lenses, focusing and defocusing the <u>radio</u> <u>waves</u>, making them appear to strengthen and weaken over a period of days, weeks or months," Dr Bannister said.

These episodes were so hard to find that researchers had given up looking for them.

But Dr Bannister and his colleagues realised they could do it with CSIRO's Compact Array.

Pointing the telescope at a quasar called PKS 1939-315 in the constellation of Sagittarius, they saw a lensing event that went on for a year.



Astronomers think the lenses are about the size of the Earth's orbit around the Sun and lie approximately 3000 light-years away - 1000 times further than the nearest star, Proxima Centauri.

Until now they knew nothing about their shape, however, the team has shown this lens could not be a solid lump or shaped like a bent sheet.

"We could be looking at a flat sheet, edge on," CSIRO team member Dr Cormac Reynolds said.

"Or we might be looking down the barrel of a hollow cylinder like a noodle, or at a spherical shell like a hazelnut."

Getting more observations will "definitely sort out the geometry," he said.

While the lensing event went on, Dr Bannister's team observed it with other radio and optical telescopes.

The optical light from the quasar didn't vary while the radio lensing was taking place. This is important, Dr Bannister said, because it means earlier optical surveys that looked for dark lumps in space couldn't have found the one his team has detected.

So what can these lenses be? One suggestion is cold clouds of <u>gas</u> that stay pulled together by the force of their own gravity. That model, worked through in detail, implies the clouds must make up a substantial fraction of the mass of our Galaxy.

Nobody knows how the invisible lenses could form. "But these structures are real, and our observations are a big step forward in determining their size and shape," Dr Bannister said.



**More information:** "Real-time detection of an extreme scattering event: Constraints on Galactic plasma lenses," <u>DOI:</u> <u>10.1126/science.aac7673</u>

## Provided by CSIRO Australia

Citation: Dark 'noodles' may lurk in the Milky Way (2016, January 21) retrieved 18 July 2024 from <u>https://phys.org/news/2016-01-dark-noodles-lurk-milky.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.