

Clarifying the role of magnetism in high-temperature superconductors

January 25 2016

A collaboration of scientists from the RIKEN SPring-8 Center, Osaka University, the Japan Atomic Energy Agency, and the Japan Synchrotron Radiation Research Institute have published research clarifying the role of magnetism in a new type of high-temperature superconductor. The research, just published as a Rapid Communication in Physical Review B, gives us a better understanding of the atomic-scale behavior of these materials. Physicists hope that, by understanding how these materials superconduct at relatively high temperature, they can eventually learn enough to make materials that superconduct close to room temperature.

It is known that the phenomenon of superconductivity—where materials conduct electricity without resistance—arises when pairs of electrons become coupled together or "paired". With traditional superconductors, this pairing arises due to vibrations of the ions in the structure. But this is not always the case: there are other types of materials, such as [cuprate superconductors](#) and a relatively new class of superconductor iron-pnictide superconductors, that was discovered by a group led by Hideo Hosono at the Tokyo Institute of Technology, where [magnetism](#) may be the pairing mechanism.

According to Alfred Baron, the leader of the Materials Dynamics Lab at RIKEN SPring-8 Center, "The question we addressed was how the [atomic vibrations](#) in the iron pnictides are affected by magnetism. This was especially interesting because atomic vibrations are understood to be driving force of the older type of low-temperature superconductors, while magnetism is considered to be the probable driving mechanism of

the new, high-temperature, superconductivity. Thus, it was in some sense, an overlap of the old with the new."

Using a technique called inelastic x-ray scattering on two beamlines of the powerful SPring-8 synchrotron facility in Harima, Japan, the group was able to measure the dynamics in specially prepared single-domain samples. Comparing their measurements to calculations then suggested that magnetic fluctuations play an important role in the atomic vibrations. Naoki Murai, the graduate student spearheading the measurement explains, "By very gently pressing the material in the correct direction we were able to observe effects due to the onset of magnetic order". Says Baron, "One of the nice things about this work is that it provides a basis for describing atomic vibrations in this whole class of materials—do calculations with magnetism and then add fluctuations".

Baron says the collaboration will continue to investigate the properties of these fascinating [materials](#), and also, more generally, the interaction of magnetism and atomic vibrations.

More information: N. Murai et al., Effect of magnetism on lattice dynamics in SrFe₂As₂ using high-resolution inelastic x-ray scattering, *Physical Review B* 93, 020301(R) (2016), DOI: 10.1103/PhysRevB.93.020301

Provided by RIKEN

Citation: Clarifying the role of magnetism in high-temperature superconductors (2016, January 25) retrieved 25 April 2024 from <https://phys.org/news/2016-01-role-magnetism-high-temperature-superconductors.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.