


# Ancient gene network helps plants adapt to their environments

February 9 2016



Timmermans and colleagues examined its function in the moss species *Physcomitrella patens*. These images show how this small RNA sensitizes the plant to auxin, an ancient plant hormone produced in response to environmental cues. These sequential images show how moss normally extends long filaments into its surroundings in response to a synthetic version of auxin called NAA. As NAA concentration increases, so does the frequency and length of these long filaments. In the bottom row, a gene (*Ppsgs3*) has been knocked out that disrupts the tasiARF pathway, resulting in a weaker response of the moss to auxin and hence far fewer and shorter protrusions. Credit: Timmermans Lab, CSHL

The only constant is change. In evolution, there are, however, some exceptions. While the enormous diversity of life suggests that organisms are constantly being refitted with new or modified parts, many of the

tools used to build these new organisms are just too useful to tinker with. For this reason they are, as scientists say, "conserved" over evolutionary time.

Professor Marja Timmermans and a team of scientists at Cold Spring Harbor Laboratory (CSHL) have discovered the purpose of one such tool that is present in both mosses and flowering plants, organisms whose common ancestor dates back 450 million years.

As they report today in *Developmental Cell*, the tool, which consists of a [gene network](#) comprising a snippet of non-coding genetic material called a small RNA and the protein it regulates, has been used over the eons to make plants more sensitive to environmental cues and facilitate robust, yet flexible, responses to those cues.

Small RNAs regulate gene activity via a mechanism called RNA interference (RNAi). One set of small RNAs called tasiARF has been highly conserved across the evolution of land plants. tasiARF regulates the genetic expression of proteins called auxin response factors (ARFs). Together, the tasiARF/ARF gene network has been found to play a role in the development of the sexual organs, roots, and leaves of [flowering plants](#). Surprisingly, this gene network is also found in mosses, which produce none of these anatomical features.

"This raises the question about what it is that tasiARF regulation really provides that is beneficial—so beneficial that it has been retained over [evolutionary time](#), and then on top of that has been used over and over and over again," says Timmermans, who is also the Alexander von Humboldt Professor at the Center for Plant Molecular Biology, University of Tübingen, Tübingen, Germany.

To reveal the secret behind the tasiARF gene network's repeated deployment, the researchers examined its function in the response of the

moss species *Physcomitrella patens* to auxin, a hormone whose effects were first studied by Charles Darwin. Auxin, which like *tasiARF* is ancient, affects plant development in a range of different ways, making regulation of the response to this hormone a key priority for plants.

The researchers discovered two benefits that *tasiARF* confers to *Physcomitrella*'s auxin response. First, *tasiARF* makes the moss more sensitive to auxin. Second, *tasiARF* makes more stable and uniform the expression of genes that are spurred upon the sensing of auxin.

While *tasiARF* affects auxin sensitivity and robustness of gene expression at the level of individual cells, Timmermans' group also discovered a unique organism-wide attribute of the small RNA. Across the system of filaments that make up the early moss plant, some cells have high levels of *tasiARF*, and are therefore more responsive to auxin, and some do not, and are therefore less responsive to the hormone.

"This is where it gets interesting for people in other fields," says Timmermans. In the presence of auxin, she says, [stem cells](#) at the tips of filaments will differentiate. It is not in the plant's best interest to allow all of the stem cells in its filament system to differentiate at once. Rather, if only a fraction of these cells differentiate, the remaining stem cells are available to respond differently in case the environment should change, providing something of a safety net.

"Once a stem cell decides to commit to becoming a certain cell type, it does so in a robust way," says Timmermans. "Across the whole organism, however, the plant can hedge its bet." Timmermans believes that gene networks like *tasiARF*/*ARF* that are regulated by small RNAs may enable similar bet-hedging strategies in other plants and animals.

It is the ability of *tasiARF* to promote a sensitive and robust auxin response, as well as to spatially regulate this response within the plant,

that Timmermans and her colleagues think made it such an indispensable tool over 450 million years of plant evolution.

"This study is important because it crosses so many disciplines," says Timmermans. "Yes, it is important to plant researchers, but it's also really adding to what we know about evolution."

**More information:** Yevgeniy Plavskin et al. Ancient trans-Acting siRNAs Confer Robustness and Sensitivity onto the Auxin Response, *Developmental Cell* (2016). [DOI: 10.1016/j.devcel.2016.01.010](https://doi.org/10.1016/j.devcel.2016.01.010)

Provided by Cold Spring Harbor Laboratory

Citation: Ancient gene network helps plants adapt to their environments (2016, February 9)  
retrieved 30 April 2024 from  
<https://phys.org/news/2016-02-ancient-gene-network-environments.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--