


## **The frigid Flying Saucer**

February 3 2016



The young star 2MASS J16281370-2431391 lies in the spectacular Rho Ophiuchi star formation region, about 400 light-years from Earth. It is surrounded by a disc of gas and dust -- such discs are called protoplanetary discs as they are the early stages in the creation of planetary systems. This particular disc is seen nearly edge-on, and its appearance in visible light pictures has led to its being nicknamed the Flying Saucer. The main image shows part of the Rho Ophiuchi region and a much enlarged close-up infrared view of the Flying Saucer from the NASA/ESA Hubble Space Telescope is shown as an insert. Credit: Digitized Sky Survey 2/NASA/ESA



The international team, led by Stephane Guilloteau at the Laboratoire d'Astrophysique de Bordeaux, France, measured the temperature of large dust grains around the young star 2MASS J16281370-2431391 in the spectacular Rho Ophiuchi star formation region, about 400 light-years from Earth.

This star is surrounded by a disc of gas and dust—such discs are called protoplanetary discs as they are the early stages in the creation of planetary systems. This particular disc is seen nearly edge-on, and its appearance in visible light pictures has led to its being nicknamed the Flying Saucer.

The astronomers used the Atacama Large Millimeter/submillimeter Array (ALMA) to observe the glow coming from carbon monoxide molecules in the 2MASS J16281370-2431391 disc. They were able to create very sharp images and found something strange—in some cases they saw a negative signal! Normally a negative signal is physically impossible, but in this case there is an explanation, which leads to a surprising conclusion.

Lead author Stephane Guilloteau takes up the story: "This disc is not observed against a black and empty night sky. Instead it's seen in silhouette in front of the glow of the Rho Ophiuchi Nebula. This diffuse glow is too extended to be detected by ALMA, but the disc absorbs it. The resulting negative signal means that parts of the disc are colder than the background. The Earth is quite literally in the shadow of the Flying Saucer!"

The team combined the ALMA measurements of the disc with observations of the background glow made with the IRAM 30-metre telescope in Spain. They derived a disc dust grain temperature of only -266 degrees Celsius (only 7 degrees above absolute zero, or 7 Kelvin) at a distance of about 15 billion kilometres from the central star. This is the



first direct measurement of the temperature of large grains (with sizes of about one millimetre) in such objects.

This temperature is much lower than the -258 to -253 degrees Celsius (15 to 20 Kelvin) that most current models predict. To resolve the discrepancy, the large <u>dust grains</u> must have different properties than those currently assumed, to allow them to cool down to such low temperatures.

"To work out the impact of this discovery on disc structure, we have to find what plausible dust properties can result in such low temperatures. We have a few ideas—for example the temperature may depend on grain size, with the bigger grains cooler than the smaller ones. But it is too early to be sure," adds co-author Emmanuel di Folco (Laboratoire d'Astrophysique de Bordeaux).

If these low dust temperatures are found to be a normal feature of protoplanetary discs this may have many consequences for understanding how they form and evolve.

For example, different dust properties will affect what happens when these particles collide, and thus their role in providing the seeds for planet formation. Whether the required change in dust properties is significant or not in this respect cannot yet be assessed.

Low dust temperatures can also have a major impact for the smaller dusty discs that are known to exist. If these discs are composed of mostly larger, but cooler, grains than is currently supposed, this would mean that these compact discs can be arbitrarily massive, so could still form giant planets comparatively close to the central star.

Further observations are needed, but it seems that the cooler dust found by ALMA may have significant consequences for the understanding of



protoplanetary discs.

This research was presented in a paper entitled "The shadow of the Flying Saucer: A very low temperature for large dust grains", by S. Guilloteau et al., published in *Astronomy & Astrophysics Letters*.

Provided by ESO

Citation: The frigid Flying Saucer (2016, February 3) retrieved 2 May 2024 from <u>https://phys.org/news/2016-02-frigid-saucer.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.