


# Scientists create ultrathin semiconductor heterostructures for new technologies

February 12 2016, by James Urton



An illustration of the strong valley exciton interactions and transport in a 2-D semiconductor heterostructure. Credit: Kyle Seyler, Pasqual Rivera

Heterostructures formed by different three-dimensional semiconductors form the foundation for modern electronic and photonic devices. Now, University of Washington scientists have successfully combined two different ultrathin semiconductors—each just one layer of atoms thick and roughly 100,000 times thinner than a human hair—to make a new two-dimensional heterostructure with potential uses in clean energy and optically-active electronics. The team, led by Boeing Distinguished Associate Professor Xiaodong Xu, announced its findings in a paper published Feb. 12 in the journal *Science*.

Senior author Xu and lead authors Kyle Seyler and Pasqual Rivera, both doctoral students in the UW physics department, synthesized and investigated the optical properties of this new type of semiconductor sandwich.

"What we're seeing here is distinct from heterostructures made of 3-D semiconductors," said Xu, who has joint appointments in the Department of Physics and the Department of Materials Science and Engineering. "We've created a system to study the special properties of these atomically thin layers and their potential to answer basic questions about physics and develop new electronic and photonic technologies."

When semiconductors absorb light, pairs of positive and negative charges can form and bind together to create so-called excitons. Scientists have long studied how these excitons behave, but when they are squeezed down to the 2-D limit in these atomically thin materials, surprising interactions can occur.

While traditional semiconductors manipulate the flow of electron charge, this device allows excitons to be preserved in "valleys," a concept from quantum mechanics similar to the spin of electrons. This is a critical step in the development of new nanoscale technologies that integrate light with electronics.

"It was already known that these ultrathin 2-D semiconductor have these unique properties that you cannot find in other 2-D or 3-D arrangements," said Xu. "But as we show here, when we put these two layers together—one on top of the other—the interface between these sheets becomes the site of even more new physical properties, which you don't see in each layer on its own or in the 3-D version."

Xu and his team wanted to create and explore the properties of a 2-D semiconductor heterostructure made up of two different layers of

material, a natural expansion of their previous studies on atomically thin junctions, as well as nanoscale lasers based on atomically thin layers of semiconductors. By studying how laser light interacts with this heterostructure, they gathered information about the physical properties at the atomically sharp interface.

"Many groups have studied the optical properties of single 2-D sheets," said Seyler. "What we do here is carefully stack one material on top of another, and then study the new properties that arise at the interface."

The team obtained two types of semiconducting crystals, tungsten diselenide (WSe<sub>2</sub>) and molybdenum diselenide (MoSe<sub>2</sub>), from collaborators at Oak Ridge National Laboratory. They used facilities developed in-house to precisely arrange two layers, one derived from each crystal, a process that took a few years to fully develop.

"But now that we know how to do it properly, we can make new ones in one or two weeks," said Xu.

Getting these devices to emit light posed a unique challenge, due to the properties of electrons in each layer.

"Once you have these two sheets of material, an essential question is how to position the two layers together," said Seyler. The electrons in each layer have unique spin and valley properties, and "how you position them—their twist angle—affects how they interact with light."

By aligning the crystal lattices, the authors could excite the heterostructure with a laser and create optically active excitons between the two layers.

"These excitons at the interface can store valley information for orders of magnitude longer than either of the layers on their own," said Rivera.

"This long lifetime allows for fascinating effects which may lead to further optical and electronic applications with valley functionality."

Now that they can efficiently make a semiconductor heterostructure out of 2-D materials, Xu and his team would like to explore a number of fascinating [physical properties](#), including how exciton behavior varies as they change angles between the layers, the quantum properties excitons between layers and electrically driven light emission.

"There's a whole industry that wants to use these 2-D semiconductors to make new electronic and [photonic devices](#)," said Xu. "So we're trying to study the fundamental [properties](#) of these new heterostructures for things like efficient laser technology, light-emitting diodes and light-harvesting devices. These will hopefully be useful for clean energy and information technology applications. It is quite exciting but there's a lot work to do."

**More information:** P. Rivera et al. Valley-polarized exciton dynamics in a 2D semiconductor heterostructure, *Science* (2016). [DOI: 10.1126/science.aac7820](#)

Provided by University of Washington

Citation: Scientists create ultrathin semiconductor heterostructures for new technologies (2016, February 12) retrieved 27 April 2024 from <https://phys.org/news/2016-02-scientists-ultrathin-semiconductor-heterostructures-technologies.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.