

# **A violent wind blown from the heart of a galaxy tells the tale of a merger**

February 4 2016

---



Figure 1: A pseudo-color image of NGC 6240 taken with Suprime-Cam at the Subaru Telescope. Blue, green, and red colors are attributed to the B-band, R-band, and H-alpha (emission line from ionized hydrogen gas) images, respectively. The giant ionized gas blown out from the galaxy is seen in red. Click the figure to jump to the larger image of the central region. Credit: Hiroshima University / NAOJ

An international team led by a researcher from Hiroshima University has succeeded in revealing the detailed structure of a massive ionized gas outflow streaming from the starburst galaxy NGC 6240 (Figure 1). The team used the Suprime-Cam mounted on the 8.2-meter Subaru Telescope on Maunakea in Hawaii.

The [ionized gas](#) the astronomers observed extends across 300,000 light-years and is carried out of the galaxy by a powerful superwind. That wind is driven by intense star-forming activity at the galactic center. The light-collecting power and high spatial resolution of Subaru Telescope made it possible to study, for the first time, the complex structure of one of the largest known superwinds being driven by starbirth – and star death.

## Starbursts and the Evolution of Galaxies

The term "starburst" indicates large-scale intensive star-forming activity, making a "starburst galaxy" one where starbirth is occurring on a grand scale. The star formation rate (SFR) of our Milky Way Galaxy is approximately one solar mass per year. By contrast, the SFRs of starburst [galaxies](#) reach ten, or even a hundred to a thousand solar masses per year.

Starburst activity is a very important part of [galaxy evolution](#). When a

starburst occurs, the intense episode star formation rapidly consumes the galaxy's interstellar gas. In addition, ultraviolet light from newborn massive stars as well as gas heating and ram pressure from supernova explosions blows much of a galaxy's gas away into intergalactic space. This galactic-scale energetic wind is called a "galactic wind" or "superwind". Its action forces interstellar gas out of the galaxy very efficiently, which accelerates the galaxy's gas-loss rate. It also chokes off [star formation](#).




Figure 2: The giant ionized gas nebula (H-alpha nebula) of NGC 6240. This image was produced by subtracting the stellar continuum light from the H-alpha image and extracting the pure (net) H-alpha emission. The complicated filamentary structure extends over hundreds of thousands light years. Credit: Hiroshima University / NAOJ

The metal-rich gas expelled from the galaxy's disk pollutes its halo as well as intergalactic space. Consequently, starburst and starburst-driven superwind significantly affect the evolution of the galaxy and the gas outside of that galaxy.

One of the mechanisms that seems to induce large-scale starburst activity is galaxy collision and merger. When two gas-rich giant spiral galaxies merge, the gravitational perturbation induced by the merging process disturbs the orbits of stars. At the same time, the gas in the galaxy disks loses its angular momentum via viscous process associated with gas mixture, and falls into the gravitational center of the merger. This creates a vast concentration of gas, which begins to coalesce, creating a starburst knot. The starburst also creates a huge amount of dust which emits strong infrared radiation as it absorbs ultraviolet light from the newly born massive stars.

## **Starburst Activity in NGC 6240**

NGC 6240 is a starburst galaxy fairly close to the Milky Way, at a distance of about 350 million light-years. Its SFR is estimated to be 25-80 times that of our galaxy (Note). It has a peculiar, disturbed morphology (Figure 1) which indicates that two spiral galaxies are merging. Due to the giant starburst at its heart as a result of the merger, NGC 6240 is very bright in infrared light being emitted from heated dust. The total infrared luminosity reaches almost a trillion times of that the Sun's.




Figure 3: (a) Right panel: Same as Figure 2. The giant ionized gas nebula (H-alpha nebula) of NGC 6240. This image was produced by subtracting the stellar continuum light from the H-alpha image and extracting the pure (net) H-alpha emission. The complicated filamentary structure extends over hundreds of thousands light years. (b) Left panel: A sketch of the H-alpha nebula of NGC 6240. Credit: Hiroshima University / NAOJ

NGC 6240 is an important object to investigate in order to understand the physical and evolutionary relationship among the processes of galaxy merger, the action of a starburst, and the phenomenon of an active galactic nucleus. Hence, it is one of the most-studied starburst galaxies in the nearby universe within 500 million light-years of the Sun.

## Observing NGC 6240 and Its Massive Outflow

The research team wanted a wide-field of NGC 6240. The Suprime-Cam optical camera was used on Subaru Telescope to zero in on the detailed

structure of the starburst-driven superwind. In addition, the team searched for important clues to understanding the starburst history of NGC 6240. They observed the galaxy using a special band-pass filter that selectively transmits the light around an emission line produced by ionized hydrogen (called the H-alpha emission line). It allowed them to study the structure of the ionized gas associated with the superwind.

## **Diving Deep into the Outflow**

Their unprecedented deep observation (long-exposure images) revealed a complex giant ionized gas nebula (called an "H-alpha nebula") surrounding NGC 6240 (Figure 2, 3a). This nebula extends out about 300,000 light-years and contains complicated structures of filaments, loops, and blobs (Figure 3b). Astronomers knew that such a large ionized [gas](#) nebula surrounds NGC 6240, but the depth of the observation significantly surpassed any previous studies and first allowed the Hiroshima team to study the some of the faintest, most detailed structure of the nebula. Large "broken bubbles" were detected in the northwestern (upper right direction of Figure 3b) and southeastern (lower left direction of Figure 3b) parts of the galaxy. These features are the evidence of a past energetic bipolar-shaped superwind that blew along the minor axis of the main galaxy disk (orthogonal to the main galactic disk).

## **Starburst-induced Superwinds Trace a Galactic Merger**

The research team performed detailed data analysis and found that NGC 6240 has experienced violent starbursts at least three times in the past and each starburst drove an energetic superwind. Those superwinds form complex structure in the H-alpha nebula. The oldest starburst started about 80 million years ago. Astronomers think that the galaxy merger



process of NGC 6240 began about a billion years ago, so this work suggests that the later stages of the merger are what excited the gigantic starbursts and subsequent superwinds. These results contribute new information to the studies of galaxy evolution and its relation to galaxy-galaxy mergers.

**More information:** Yoshida et al., 2016, "Giant H-alpha Nebula Surrounding the Starburst Merger NGC 6240" *Astrophysical Journal*

Provided by Subaru Telescope

Citation: A violent wind blown from the heart of a galaxy tells the tale of a merger (2016, February 4) retrieved 26 April 2024 from <https://phys.org/news/2016-02-violent-blown-heart-galaxy-tale.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.