

Fires are increasing in warming world, but a new model could help us predict them

February 10 2016, by Ritaban Dutta, Csiro



Credit: AI-generated image (disclaimer)

Over the past decade, the frequency of bushfires in Australia <u>has</u> <u>increased</u>. The <u>Forest Fire Danger Index</u> - which measures the frequency and severity of the weather most conducive to fire - has increased dramatically since the 1970s, and particularly in the 1990s and 2000s. This – along with a great deal of other evidence – indicates that a major



change in the climate is looming.

In a study published today in *Royal Society Open Science*, we show that <u>bushfire</u> frequency has increased by 40% over the past five years.

This is part of our work to better predict bushfires using weather information. We developed a model that can predict bushfires over a week-long period with an accuracy of 91%. While we can't yet use this to predict fires definitively, it is a big step forward in planning for and predicting fires.


There is an urgent need for scientific research that can contribute to Australia's future bushfire preparedness. The knowledge gained from this research could help save lives and property, underpin effective building codes and help land-management decisions.

Fire forecasting gets smart

We wanted to find bushfire "hotspots" – places where a bushfire might happen in the future based on recent weather patterns. They don't show that a bushfire will definitely happen, and similarly we need to be careful that people don't assume a fire won't happen if there is no hotspot.

To do this we developed a model that combined data on fires (from satellite images) with data on climate, including soil moisture, dry fuel load, wind speed, temperature and humidity.





Map showing predicted hotspots (green) and real fires (purple). Red dots show fires not predicted by the model. Credit: Ritaban Dutta

We are using a <u>machine learning</u> approach to model future bushfire hotspots - in short, a form of artificial intelligence that allows the model to learn on its own. Our model used other forms of learning to produce maps of bushfire hotspots.

Predicting hotspots

We tested our model by looking at bushfires retrospectively. Our predictions achieved 91% global accuracy. The analysis also indicates that, on a week-by-week basis, the frequency of Australian bushfires, particularly during summer months, has increased by 40% over the past five years.

We conducted extensive research in Australian bushfire history, to make sure the data on the conditions and behaviour of 36 major fires were correct.



Two types of Australian vegetation are prone to fire: grasslands and forests. We found that our model could predict both.

Fighting fires

Although new technologies are being developed to manage the growing number of unplanned bushfires in Australia, we do not have, and are never likely to have, a way to avert natural fire disasters.

The recent history of Australian bushfires indicates that the most effective way to save lives is early planned evacuation, combined with timely advice and alerts to the people potentially at risk.

Early estimation of the likely frequency of future bushfires, and accurate hot-spot estimation of the locations most likely to be affected could provide great support to land managers, and assist in reducing the damage caused.

But as yet, our system exists only as a research <u>model</u>. Its impact and effectiveness could only be fully evaluated if a real system were implemented as a decision support tool for land managers and emergency services, to accurately forecast the most likely bushfire hot spots, at a detailed and fine-grained resolution and a timely scale.

More information: Big data integration shows Australian bush-fire frequency is increasing significantly, *Royal Society Open Science*, <u>rsos.royalsocietypublishing.or</u> ... /10.1098/rsos.150241

This article was originally published on <u>The Conversation</u>. *Read the* <u>original article</u>.



Source: The Conversation

Citation: Fires are increasing in warming world, but a new model could help us predict them (2016, February 10) retrieved 25 April 2024 from <u>https://phys.org/news/2016-02-world-predictthem.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.