


# Two new zoantharian species found on eunicid worms in the dark in the Indo-Pacific ocean

February 11 2016

---



One of the new zoantharian species, *Epizoanthus inazuma*. Credit: Hiroki Kise

While studying the abundant, yet poorly known fauna of the zoantharian *Epizoanthus* genus in the Indo-Pacific Ocean, Japanese graduate student Hiroki Kise and Dr. James Davis Reimer, both affiliated with University of the Ryukyus, Japan, focused on examining the species living on eunicid worm tubes. These *Epizoanthus* species form colonies and are only found living on the outside surfaces of tubes in which the eunicid worms live. Although these zoantharians often live in areas that are deeper than the range of SCUBA divers and tend to be indistinguishable on the outside, the present research, published in the open-access journal *ZooKeys*, reports the discovery of two new species.

Both [new species](#) have been found in low-light environments. While one of the species, scientifically named *Epizoanthus inazuma*, showed preference for either coral reef slopes, reef floors, or the sides of their overhangs in Okinawa, the other one, *E. beriber* was seen to tolerate cave environments exclusively, and is found in Palau and Papua New Guinea. To recognise them as new species, the researchers turned to molecular analyses combined and compared to morphological data. Thus, this discovery presents yet another example of the utility of molecular methods as an effective tool in taxonomic and biodiversity research.

Forming colonies resembling a classic lightning-bolt shape, the new coral species *Epizoanthus inazuma* has reasonably been given a name meaning 'lightning' in Japanese. The second new species is also named in a reference to its lifestyle as it bears the name of the local Palauan folklore character Beriber, who lived in a cave.

The researchers believe that it is highly likely that there are other undescribed species in coral reefs. There is even greater likelihood that such are currently hidden in underwater cave habitats. The authors do not exclude the possibility that these new zoantharian species may be distributed across other locations in the Pacific, where they could easily be mistaken for other closely related [species](#).



A colony of the new zoantharian *Epizoanthus beriber* species. Credit: Hiroki Kise

**More information:** Hiroki Kise et al. Unexpected diversity and a new species of *Epizoanthus* (Anthozoa, Hexacorallia) attached to eunicid worm tubes from the Pacific Ocean, *ZooKeys* (2016). [DOI: 10.3897/zookeys.562.6181](https://doi.org/10.3897/zookeys.562.6181)

Provided by Pensoft Publishers

Citation: Two new zoantharian species found on eunicid worms in the dark in the Indo-Pacific ocean (2016, February 11) retrieved 16 August 2024 from

<https://phys.org/news/2016-02-zoantharian-species-eunicid-worms-dark.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.