

Continuing the search for gravitational waves

March 29 2016, by Peter Garnavich



LBT (Large Binocular Telescope)

In February, the LIGO Scientific Collaboration announced it had detected gravitational waves for the first time, confirming the last prediction of Albert Einstein's theory of relativity. Somewhat overlooked in the excitement that followed is the fact that scientists don't know the exact location the waves were coming from. University of Notre Dame astronomer Peter Garnavich is leading a group of researchers who are hoping to more precisely locate where future gravitational waves originate.

Garnavich and the group are using the LBT (Large Binocular Telescope) in southeastern Arizona to search for visible light emission from the event generating the gravitational waves. Notre Dame owns a share of the LBT project, which consists of two 8.4-meter mirrors and is the

world's largest telescope on a single mounting. The team is searching the sky to find the light emitted from supernovae and gamma ray bursts, and now gravitational wave transients.

"Gravitational wave transients are like earthquakes," Garnavich said. "Earthquakes make waves that propagate around and through the earth. A truck hitting a pothole will send out some vibrations, but they won't make it very far. A big earthquake can send waves out that can be felt hundreds of miles away. So these two [black holes](#) merging are like a huge earthquake, but instead of the earth shaking, it is space-time vibrating. Advanced LIGO is like a seismograph for space-time."

Garnavich notes that the gravitational waves that the LIGO group detected emanating from two black holes merging were somewhat of a surprise.

"These LIGO black holes are bigger than the stellar black holes we know about in the Milky Way, but much smaller than the massive black holes in the centers of galaxies. This may be as big a discovery as the gravitational waves themselves," Garnavich said. "Black hole mergers are not expected to produce a lot of electromagnetic radiation, and the prevailing consensus was that gravitational waves would be detected from colliding neutron stars, an event that should generate some light."

Garnavich notes that the effort to more precisely locate gravitational waves is akin to searching for a needle in a haystack. The swath of space that LIGO's research identified as a possible location of the event can contain millions of stars and galaxies.

"We hunt for things that snap, crackle or pop in their light output, and that narrows candidates down to maybe 100 across the sky," he said

Garnavich's group is hoping to use the LBT and other observational

surveys to identify promising candidates and eliminate not-so-promising ones.

"I hope this LIGO event was the first of many so we can combine [gravitational waves](#) and [electromagnetic waves](#) to figure out what are creating these exotic events," he said.

Given past experience, a breakthrough like the LIGO discovery opens the door to even more discoveries. The search Garnavich and his group is undertaking has the potential to provide a new understanding of how the universe works.

Provided by University of Notre Dame

Citation: Continuing the search for gravitational waves (2016, March 29) retrieved 20 April 2024 from <https://phys.org/news/2016-03-gravitational.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.