


Eating green could be in your genes

March 29 2016



Global frequency pattern of an allele adaptive to vegetarian diets. The allele has the highest frequency in Indians, who have traditionally relied heavily on plant-based diets. Credit: Kaixiong Ye, Cornell University/Map from outline-world-map.com

Could there be a vegetarian gene? Cornell University researchers have found evidence of a genetic variation - called an allele - that has evolved in populations that have historically favored vegetarian diets, such as in India, Africa and parts of East Asia. They also discovered a different version of this gene adapted to a marine diet discovered among the Inuit in Greenland, who mainly consume seafood.

The vegetarian allele evolved in populations that have eaten a plant-based diet over hundreds of generations. The adaptation allows these people to efficiently process omega-3 and omega-6 [fatty acids](#) and convert them into compounds essential for early brain development and if they stray from a balanced omega-6 to omega-3 diet, it may make people more susceptible to inflammation, and by association, increased risk of heart disease and colon cancer.

In Inuit populations of Greenland, the researchers uncovered that a previously identified adaptation is opposite to the one found in long-standing vegetarian populations: While the vegetarian allele has an insertion of 22 bases (a base is a building block of DNA) within the gene, this insertion was found to be deleted in the seafood allele.

"The opposite allele is likely driving adaptation in Inuit," said Kaixiong Ye, co-lead author of the paper appearing March 29 in the journal *Molecular Biology and Evolution*. Ye is a postdoctoral researcher in the lab of Alon Keinan, associate professor of biological statistics and computational biology, and the paper's co-senior author.

"Our study is the first to connect an insertion allele with vegetarian diets, and the deletion allele with a marine diet," Ye said.

"It is the most interesting example of local adaptation that I have been fortunate to help study," said Keinan. "Several studies have pointed to adaptation in this region of the genome. Our analyses combine to show that the adaptation is driven by an insertion of a small piece of DNA that we know its function. Moreover, when it reached the Greenlandic Inuit, with their marine-based diet rich in omega-3, it might have become detrimental."

FADS1 and FADS2 are enzymes that are essential for converting omega-3 and [omega-6 fatty acids](#) into downstream products needed for

brain development and controlling inflammation. Meat and seafood eaters have less need for increased FADS1 and FADS2 enzymes to get proper nutrition because their omega-3 and [omega-6](#) fatty acid conversion process is simpler and requires fewer steps.

This study is based on previous work by co-senior author Tom Brenna, professor of human nutrition and of chemistry at Cornell University, who showed the insertion can regulate the expression of FADS1 and FADS2 and hypothesized it could be an adaptation in vegetarian populations.

Ye, Keinan and colleagues analyzed frequencies of the vegetarian allele in 234 primarily vegetarian Indians and 311 U.S. individuals and found the vegetarian allele in 68 percent of the Indians and in just 18 percent of Americans. Analysis using data from the 1,000 Genomes Project similarly found the vegetarian allele in 70 percent of South Asians, 53 percent of Africans, 29 percent of East Asians and 17 percent of Europeans.

"Northern Europeans have a long history of drinking milk and they absorbed enough end products from milk for long-chain fatty acid metabolism so they don't have to increase capacity to synthesize those fatty acids from precursors," said Ye.

"One implication from our study is that we can use this genomic information to try to tailor our diet so it is matched to our genome, which is called personalized nutrition," he added.

The researchers are not sure yet when the adaptation first occurred, as analyses of chimpanzee or orangutan genomes did not uncover the vegetarian allele. But there is evidence for the allele in early hominid Neanderthal and Denisovan genomes.

"It is possible that in the history of human evolution, when people migrated to different environments, sometimes they ate a plant-based diet and sometimes they ate a marine-based [diet](#), and in different time periods these different alleles were adaptive," meaning the alleles have a tendency to evolve under dietary pressures, Ye said.

Provided by Cornell University

Citation: Eating green could be in your genes (2016, March 29) retrieved 9 April 2024 from <https://phys.org/news/2016-03-green-genes.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--