

New insight into enzyme evolution

March 3 2016

How enzymes - the biological proteins that act as catalysts and help complex reactions occur - are 'tuned' to work at a particular temperature is described in new research from groups in New Zealand and the UK, including the University of Bristol.

Professor Vic Arcus (University of Waikato) and colleagues, including Bristol's Professor Adrian Mulholland and Dr Marc van der Kamp, showed that the heat capacity of enzymes changes during a reaction as the enzymes tighten up. Exactly how much the enzymes tighten up is the critical factor in determining the temperature at which they work best. These findings could provide a route to designing better biocatalysts for use in [chemical reactions](#) in industrial processes, such as the production of drugs.

Enzymes have an optimum temperature at which they are most catalytically active. Above that temperature, they become less active. Previously, it was thought that this was because enzymes unfolded (lost their functional shape) at higher temperatures, but actually they typically become less active at higher temperatures even though they maintain their functional shape.

So what makes them less active? And what is it that causes enzymes from different organisms to have different catalytic activities at the same temperature? Enzymes from organisms that live at normal temperatures are not very active at low temperatures, while cold-adapted enzymes are active in the cold - why, when they have very similar structures?

The new research, published as a 'New Concept' in *Biochemistry* (and selected for the American Chemical Society (ACS) Editors' Choice), shows that a basic physical property - the heat capacity - explains and predicts the temperature dependence of enzymes. The heat capacity of a substance is the amount of heat required to raise its temperature by one degree. For enzymes, the heat capacity changes during the reaction and this change is 'tuned' to give the optimal temperature.

Professor Mulholland said: "Our theory - macromolecular rate theory, (MMRT) - applies to all enzymes, and so will have a critical role in predicting metabolic activity as a function of temperature.

"We also expect to see characteristics of MMRT at the level of cells, whole organisms and even ecosystems. This means that it is important in understanding and predicting the response of biological systems to temperature changes, for example, how ecosystems will respond to temperature changes associated with climate change."

The theory also explains why enzymes are so big (the more 'difficult' the chemistry to catalyse, the bigger the [enzyme](#)). It also hints at why proteins were eventually preferred by evolution over nucleic acids as catalysts in biology: proteins offer much more ability to 'tune' dynamics and their response to chemical reactions.

Provided by University of Bristol

Citation: New insight into enzyme evolution (2016, March 3) retrieved 3 May 2024 from <https://phys.org/news/2016-03-insight-enzyme-evolution.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--