

Newly discovered bacteria can eat plastic bottles

March 11 2016, by Deborah Netburn, Los Angeles Times



Credit: Charles Rondeau/ public domain

A team of Japanese scientists has found a species of bacteria that eats

the type of plastic found in most disposable water bottles.

The discovery, published Thursday in the journal *Science*, could lead to new methods to manage the more than 50 million tons of this particular type of plastic produced globally each year.

The plastic found in [water bottles](#) is known as polyethylene terephthalate, or PET. It is also found in polyester clothing, frozen-dinner trays and blister packaging.

"If you walk down the aisle in Wal-Mart you're seeing a lot of PET," said Tracy Mincer, who studies plastics in the ocean at the Woods Hole Oceanographic Institution in Massachusetts.

Part of the appeal of PET is that it is lightweight, colorless and strong. However, it has also been notoriously resistant to being broken down by [microbes](#)-what experts call "biodegradation."

Previous studies had found a few species of fungi can grow on PET, but until now, no one had found any microbes that can eat it.

To find the plastic-eating bacterium described in the study, the Japanese research team from Kyoto Institute of Technology and Keio University collected 250 PET-contaminated samples including sediment, soil and wastewater from a plastic bottle recycling site.

Next they screened the microbes living on the samples to see whether any of them were eating the PET and using it to grow. They originally found a consortium of bugs that appeared to break down a PET film, but they eventually discovered that just one of [bacteria species](#) was responsible for the PET degradation. They named it *Ideonella sakainesis*.

Further tests in the lab revealed that it used two enzymes to break down

the PET. After adhering to the PET surface, the bacteria secretes one enzyme onto the PET to generate an intermediate chemical. That chemical is then taken up by the cell, where another enzyme breaks it down even further, providing the bacteria with carbon and energy to grow.

The researchers report that a community of *Ideonella sakaiensis* working this way could break down a thin film of PET over the course of six weeks if the temperature were held at a steady 86 degrees Fahrenheit.

Mincer said the study was impressive and did a good job showing that these organisms were eating the plastic pretty well. However, he said it was not immediately clear whether or not it would help keep plastics out of the ocean, for example.

"When I think it through, I don't really know where it gets us," he said. "I don't see how microbes degrading plastics is any better than putting [plastic](#) bottles in a recycling bin so they can be melted down to make new ones."

He added that the research could make it easier to identify other microbes that might have similar PET-degrading capabilities.

"This process could be quite common," he said. "Now that we know what we are looking for, we may see these microbes in many areas around the world."

More information: U. T. Bornscheuer. Feeding on plastic, *Science* (2016). [DOI: 10.1126/science.aaf2853](https://doi.org/10.1126/science.aaf2853)

S. Yoshida et al. A bacterium that degrades and assimilates poly(ethylene terephthalate), *Science* (2016). [DOI: 10.1126/science.aad6359](https://doi.org/10.1126/science.aad6359)

©2016 Los Angeles Times
Distributed by Tribune Content Agency, LLC.

Citation: Newly discovered bacteria can eat plastic bottles (2016, March 11) retrieved 10 April 2024 from <https://phys.org/news/2016-03-newly-bacteria-plastic-bottles.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.