


Second quantum revolution a reality with chip-based atomic physics

March 30 2016



The image shows a quartz surface above the electrodes used to trap atoms. The color map on the surface shows the electric field amplitude. Credit: University of Oklahoma

A University of Oklahoma-led team of physicists believes chip-based atomic physics holds promise to make the second quantum revolution—the engineering of quantum matter with arbitrary precision—a reality. With recent technological advances in fabrication and trapping, hybrid quantum systems are emerging as ideal platforms for a diverse range of studies in quantum control, quantum simulation

and computing.

James P. Shaffer, professor in the Homer L. Dodge Department of Physics and Astronomy, OU College of Arts and Sciences; Jon Sedlacek, OU graduate student; and a team from the University of Nevada, Western Washington University, The United States Naval Academy, Sandia National Laboratories and Harvard-Smithsonian Center for Astrophysics, have published research important for integrating Rydberg atoms into hybrid quantum systems and the fundamental study of atom-[surface](#) interactions, as well as applications for electrons bound to a 2D surface.

"A convenient surface for application in hybrid quantum systems is quartz because of its extensive use in the semiconductor and optics industries," Sedlacek said. "The surface has been the subject of recent interest as a result of its stability and low surface energy. Mitigating electric fields near 'trapping' surfaces is the holy grail for realizing hybrid [quantum systems](#)," added Hossein Sadeghpour, director of the Institute for Theoretical Atomic Molecular and Optical Physics, Harvard-Smithsonian Center for Astrophysics.

In this work, Shaffer finds ionized electrons from Rydberg atoms excited near the quartz surface form a 2D layer of electrons above the surface, canceling the electric field produced by rubidium surface adsorbates. The system is similar to electron trapping in a 2D gas on superfluid liquid helium. The binding of electrons to the surface substantially reduces the electric field above the surface.

"Our results show that binding is due to the image potential of the electron inside the quartz," said Shaffer. "The electron can't diffuse into the quartz because the rubidium adsorbates make the surface have a negative electron affinity. The approach is a promising pathway for coupling Rydberg atoms to surfaces as well as for using surfaces close to

atomic and ionic samples."

Provided by University of Oklahoma

Citation: Second quantum revolution a reality with chip-based atomic physics (2016, March 30)
retrieved 18 April 2024 from

<https://phys.org/news/2016-03-quantum-revolution-reality-chip-based-atomic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.