

## **Down the rabbit hole: How electrons travel through exotic new material**

March 10 2016



New research from Princeton University demonstrates the bizarre movement of electrons through a novel material called a Weyl semi-metal. The image shows a schematic of the connections at special values of electron momentum, which come in pairs and are called Weyl points (red and blue dots). An electron that leaves the surface on a red point can return through its partner blue point, and



vice versa. This bizarre behavior is due to the 'topological'connections through the bulk of the material. Credit: Yazdani et al., Princeton University.

Researchers at Princeton University have observed a bizarre behavior in a strange new crystal that could hold the key for future electronic technologies. Unlike most materials in which electrons travel on the surface, in these new materials the electrons sink into the depths of the crystal through special conductive channels.

"It is like these electrons go down a rabbit hole and show up on the opposite surface," said Ali Yazdani, the Class of 1909 Professor of Physics. "You don't find anything else like this in other materials."

The research was published in the journal Science.

Yazdani and his colleagues discovered the odd behavior while studying electrons in a crystal made of layers of tantalum and arsenic. The material, called a Weyl semi-metal, behaves both like a metal, which conducts electrons, and an insulator, which blocks them. A better understanding of these and other "topological" materials someday could lead to new, faster electronic devices.

The team's experimental results suggest that the surface electrons plunge into the crystal only when traveling at a certain speed and direction of travel called the Weyl momentum, said Yazdani. "It is as if you have an electron on one surface, and it is cruising along, and when it hits some special value of momentum, it sinks into the crystal and appears on the opposite surface," he said.





Three-dimensional image using scanning tunneling electron microscopy of electrons on the surface of a Weyl semi-metal, a kind of crystal with unusual conducting and insulating properties. Credit: Image credit: Yazdani et al., Princeton University.

These special values of momentum, also called Weyl points, can be thought of as portals where the electrons can depart from the surface and be conducted to the opposing surface. The theory predicts that the points come in pairs, so that a departing electron will make the return trip through the partner point.

The team decided to explore the behavior of these electrons following research, published in *Science* last year by another Princeton team and



separately by two independent groups, revealing that electrons in Weyl semi-metals are quite unusual. For example, their experiments implied that while most surface electrons create a wave pattern that resembles the spreading rings that ripple out when a stone is thrown into a pond, the <u>surface electrons</u> in the <u>new materials</u> should make only a half circle, earning them the name "Fermi arcs."

To get a more direct look at the patterns of electron flow in Weyl semimetals, postdoctoral researcher Hiroyuki Inoue and graduate student András Gyenis in Yazdani's lab, with help from graduate student Seong Woo Oh, used a highly sensitive instrument called a scanning tunneling microscope, one of the few tools that can observe electron waves on a crystal surface. They obtained the tantalum arsenide crystals from graduate student Shan Jiang and assistant professor Ni Ni at the University of California-Los Angeles.



Comparison of experiment (left) to theory (right): The image on the left shows patterns of waves imaged with scanning tunneling microscopy, revealing all the



possible ways in which electrons can interfere with each other on the surface. This pattern is dictated by the connection of the surface electrons with the interior of the crystal, which is determined by the Weyl momentum of electrons, the special momentum when the electrons sink easily through the sample. The observed pattern closely matches the pattern predicted by theoretical calculations (right). Credit: Image credit: Yazdani et al., Princeton University.

The results were puzzling. "Some of the interference patterns that we expected to see were missing," Yazdani said.

To help explain the phenomenon, Yazdani consulted B. Andrei Bernevig, associate professor of physics at Princeton, who has expertise in the theory of topological materials and whose group was involved in the first predictions of Weyl semi-metals in a 2015 paper published in *Physical Review X*.

Bernevig, with help from postdoctoral researchers Jian Li and Zhijun Wang, realized that the observed pattern made sense if the <u>electrons</u> in these unusual materials were sinking into the bulk of the crystal. "Nobody had predicted that there would be signals of this type of transport from a scanning tunneling microscope, so it came as a bit of a surprise," said Bernevig.

The next step, said Bernevig, is to look for the behavior in other crystals.

**More information:** "Quasiparticle interference of the Fermi arcs and surface-bulk connectivity of a Weyl semimetal," Hiroyuki Inoue, András Gyenis, Zhijun Wang, Jian Li, Seong Woo Oh, Shan Jiang, Ni Ni, B. Andrei Bernevig, and Ali Yazdani, *Science*, March 10, 2016. <u>science.sciencemag.org/cgi/doi ... 1126/science.aad8766</u>



## Provided by Princeton University

Citation: Down the rabbit hole: How electrons travel through exotic new material (2016, March 10) retrieved 15 May 2024 from https://phys.org/news/2016-03-rabbit-hole-electrons-exotic-material.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.