

Test aims to identify shale gas hazard in groundwater

April 22 2016

A test has been developed to check for contamination of shallow groundwater from unconventional gas extraction techniques, such as fracking.

The development could help monitor the safety of shale gas and coal bed [methane](#) extraction, which has sparked debate because of the perceived risks of water contamination.

Methods used for [shale gas extraction](#) include hydraulic fracturing, commonly known as fracking, in which shale rocks below ground are split with high-pressure fluids to release gas that is recovered for fuel. Coal bed methane is extracted from deep coal seams by drilling into the coal to reduce the pressure and release gas.

The need for such a test was highlighted following allegations in the US that, following fracking operations, drinking water had become contaminated with large amounts of methane from deep below ground.

Groundwater often contains methane gas from shallow natural sources, which is harmless in small quantities. The researchers have developed a new way to fingerprint [methane gas](#) by identifying tiny traces of inactive natural gases, known as noble gases. These fingerprints vary depending on the origin and depth of the methane, and enable scientists to pinpoint its source.

The researchers, from the Universities of Edinburgh and Glasgow and

the Scottish Universities Environmental Research Centre, have recorded these unique fingerprints in a number of exploratory [shale gas](#) and [coal bed methane](#) wells from around the UK. They are presenting these findings today (Friday) at the European Geosciences Union Conference in Vienna.

The fingerprint analysis can be used to determine the origin of methane at exploration sites. If levels of methane in groundwater are found to have changed following exploration activity, and the gas is traced to exploration or extraction activity, appropriate action can be taken.

Scientists developed the test by adapting a technique used for monitoring potential leaks of [carbon dioxide gas](#) from storage sites deep underground.

Dr Stuart Gilfillan, of the University of Edinburgh's School of GeoSciences, who led the project, said: "Creating this fingerprint test will enable [gas](#) exploration and extraction to be carried out responsibly and should help address public concerns over this technology. It is important that careful monitoring of methane levels in nearby waters is carried out when commercial extraction begins."

Provided by University of Edinburgh

Citation: Test aims to identify shale gas hazard in groundwater (2016, April 22) retrieved 2 May 2024 from <https://phys.org/news/2016-04-aims-shale-gas-hazard-groundwater.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--