

New data analysis technique distinguishes active from passive fluctuations inside cells

April 28 2016, by Jennifer Chu



Any system in thermodynamic equilibrium is known to satisfy perfectly balanced forward and backward transitions between any two states. It is therefore impossible, for instance, to drive a windmill to do work by using only thermal fluctuations. The new research shows how living systems violate this principle. Thus, even stochastic fluctuations in such living systems could be used to drive such a small-scale windmill. Credit: C. Hohmann and M. Leunissen



Inside every living cell, internal structures are continuously moving about. Under a microscope, organelles such as the nucleus, mitochondria, transport vesicles, or even external flagella wobble and twitch. This may happen spontaneously as these tiny structures are passively jostled inside a cell. But that's not necessarily all there is to it. Often a cell invests extra energy into these motions to enhance cell functions in ways we don't yet understand.

At the microscopic scale, particles in a fluid or a gas can move about in response to bombardment by the surrounding molecules. Such passive, thermally-induced motions are often very hard to distinguish from actively driven movements, and it can be impossible to tell just by looking whether particular motions inside a cell are simply thermal or pushed by some extra input of energy.

Now scientists at MIT, the University of Göttingen, Ludwig Maximilians University of Munich, the Free University Amsterdam, and Yale University have developed a noninvasive data analysis technique that can discern whether an object's random motion is actively or thermally driven. After tracking the conformations or locations that a structure within the cell or a particle passes through as it moves, and observing how the particle transitions back and forth between such states, the researchers apply a fundamental principle of statistical physics to determine whether the random motions are active or thermal.

Nikta Fakhri, assistant professor in MIT's Department of Physics, says the results will help scientists to uncover "hidden" active processes that drive a cell's constituents to move in seemingly random ways.

"We want to see if particular dynamics in living systems—be those cells or tissues or whole organisms—that look at first glance like random thermal motion are indeed actively driven," says Fakhri, who is a first coauthor on the paper. "This is important because there must be a vital



function connected with the process if the cell spends energy on it. Our work provides a practical experimental method to identify active, nonequilibrium processes in observations of biological systems."

Fakhri and her colleagues have published their results today in the journal *Science*.

Shaking things up

Observations of the movements of microscopic particles date back to 1827, when Scottish botanist Robert Brown was looking through a microscope at pollen grains in water. He noticed that the grains contained tiny particles that jiggled vigorously, and at first glance believed that their motion meant the particles were somehow alive and moving on their own. Eventually, Brown retired this theory when he observed the same jiggling with inert particles from rocks, realizing that just seeing a particle moving doesn't necessarily mean it is alive.

Scientists now know that such particles are being pushed around by even smaller water molecules, which themselves are jiggling. At elevated temperatures, water molecules possess kinetic energy and can remain forever restless. At microscopic scales, these tiny molecules and atoms can "bombard" other much larger particles. This phenomenon is now known as Brownian motion. Scientists today are so used to seeing Brownian motion that they often assume that a particle moving in a random fashion is likely to be at thermal equilibrium—a state in which a system is not dissipating any energy, and therefore inanimate.

In living cells, of course, many organelles or particles tend to move in a "ballistic" fashion, traveling across some distance, with a clearly directed or oscillatory trajectory. These types of motions, Fakhri says, represent an animated state that must be out of equilibrium, requiring a system to expend energy. What is interesting, however, are situations where



particles jiggle randomly and seem to be in equilibrium, but in reality move actively.

Frame by frame

Fakhri and her colleagues set out to develop a <u>statistical physics</u> technique that would enable them to tell, just by imaging a particle, whether its random motions are thermal or active in nature.

Using video microscopy, they studied, frame by frame, the oscillatory motion of the flagellum of a Chlamydomonas algae. They deconstructed the backbone of the flagellum into a series of shapes, thus creating a phase space of the states the flagellum passes through as it completes an oscillatory cycle. They then counted the transitions between states. In thermal equilibrium, the back-and-forth transitions between all states must be balanced. However, they observed a clear imbalance in these transitions, confirming the already known fact that the flagellum expends energy for this active oscillatory motion.

Next, they analyzed the motions of a kidney cell's cilium—an antennalike appendage that at first glance appears to be jiggling back and forth passively. By tracking the cilium's orientation and curvature, and counting transitions between states, they observed a slight imbalance in the transitions, pointing to an unexpected active process that drives the cilium, despite its passive appearance.

Fakhri says the new method will help scientists to uncover new ways in which cells dissipate energy—which, ultimately, is the key to sustaining life. After all, as the Austrian physicist Erwin Schrödinger noted, "Living matter evades the decay to thermal equilibrium."

"Almost a century later, through the work of this collaboration, we can now go further and find out how <u>cells</u> budget their energy and



particularly why they often expend a lot of energy to create these apparently <u>random motions</u>," Fakhri says.

More information: "Broken detailed balance at mesoscopic scales in active biological systems" *Science*, <u>DOI: 10.1126/science.aac8167</u>

Provided by Massachusetts Institute of Technology

Citation: New data analysis technique distinguishes active from passive fluctuations inside cells (2016, April 28) retrieved 2 May 2024 from <u>https://phys.org/news/2016-04-analysis-technique-distinguishes-passive-fluctuations.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.