

Study highlights processes shaping species coexistence and potential impacts of deforestation

April 12 2016, by Marilyn Siderwicz



Bialoweiza Forest is one of the last and largest remaining parts of the immense primeval forest that once stretched across the European Plain. Credit: Adam Wajrak

Białowieża Forest in Poland has been in the news lately due to the major destruction it is facing from potential deforestation. Preserved through the ages—first used as a royal game reserve, then later protected through national park initiatives and management—a movement is now underway to tap the forest for increased logging.


New research by Department of Civil and Environmental Engineering (CEE) Assistant Professor Serguei Saavedra, working with a team of European ecology researchers, highlights some of the major processes shaping the [forest](#)'s delicate [species](#) coexistence which could be dramatically altered through deforestation. Their paper is the cover story of this month's issue of *Ecology*.

The team's results imply that large anthropogenic effects, such as deforestation, can have significant negative effects on the balance of the entire ecosystem, because predator-prey interactions are closely coupled to the environment.

Only approximately 550 square miles, the forest supports thousands of different species including hundreds of bison; large carnivores such as wolf and lynx; birds; and lichens, fungi, and microbes that thrive on woodsy decomposition.

Developing strategies for sustainability

Białowieża Forest represents the last primeval temperate forest in lowland Europe. The complexity of the forest structure and the strong seasonality generate significant variations in the environmental conditions throughout the year. Saavedra and his colleagues showed that seasonal interaction turnover among species helps minimize the impact of these variations.



The top and bottom figures correspond, respectively, to a subsample of the summer and winter predator-prey interaction networks in Poland's Bialowieza Forest. In each interaction network, predators are at the top and prey at the bottom. Species in black and color correspond, respectively, to permanent and seasonal species. Black and colored lines correspond to interactions among permanent species that are present the entire year (permanent interactions) and in one season only, respectively. Dashed lines represent interactions either between permanent and seasonal species, or among seasonal species only.

"Many of the species interactions in ecological communities are not permanent, but are characterized by temporal changes," Saavedra explains. "Understanding how and why these interactions change can shed new light on the processes shaping the delicate coexistence of large ecological systems, and lead to more efficient strategies for their sustainability."

Associate Professor Nuria Selva, one of Saavedra's co-authors from the Polish Academy of Sciences, adds that Białowieża Forest is the best preserved lowland forest of the temperate zone and therefore was an excellent area to research.

"Places like this which are changed very little by people are rare, particularly in Europe," she says, "but they give a unique opportunity to observe natural processes and how the natural world is organized. Our study demonstrates that the temporal changes in how predators and their prey interact are crucial for the persistence of the whole community. This is often hard to understand by nature managers who tend to intervene as soon as they see a big change happening."

The team investigated the extent to which changes of species composition impact the likelihood of species coexistence in the highly seasonal forest, and the extent to which seasonal changes of predator diet modulate the expected impact. This likelihood was estimated by extending recent developments in the field of quantitative ecology.

The scientists found a significant species turnover in the Białowieża community between summer and winter. Importantly, they demonstrated that active changes in predators' diet maintain similar coexistence conditions through the year despite species turnover. Paradoxically, the results suggest that short-term dynamics can be coupled to the long-term persistence of ecological systems.

"Humans are altering many facets of the environment at a speed and extent that is unprecedented in the Earth's recent history," says Canterbury University Professor Jason M. Tylianakis, a terrestrial ecosystems conservation researcher who was not involved in this research. "The complexity of ecosystems, with hundreds to thousands of species interacting with each other at a given location, makes it difficult to understand, let alone predict, their responses to this change. This

research provides a key step forward by using a newly-developed analytical approach to show the importance of a few key predator-prey interactions for making the whole ecosystem persist. In short, the work of Saavedra and colleagues adds another weapon to the arsenal in the fight against species extinctions."

More information: Serguei Saavedra et al. Seasonal species interactions minimize the impact of species turnover on the likelihood of community persistence, *Ecology* (2015). [DOI: 10.1890/15-1013.1](https://doi.org/10.1890/15-1013.1)

This story is republished courtesy of MIT News (web.mit.edu/newsoffice/), a popular site that covers news about MIT research, innovation and teaching.

Provided by Massachusetts Institute of Technology

Citation: Study highlights processes shaping species coexistence and potential impacts of deforestation (2016, April 12) retrieved 25 April 2024 from <https://phys.org/news/2016-04-highlights-species-coexistence-potential-impacts.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--