


Mystery solved: Traits identified for why certain chemicals reach toxic levels in food webs

April 20 2016



This model provides a new global tool for screening existing and new organic chemicals for their biomagnification potential. Hot colors (red, orange and yellow) indicate a high probability of biomagnification and cool colors (greens, blues) indicate a low probability of biomagnification. Credit: USGS

Researchers have figured out what makes certain chemicals accumulate to toxic levels in aquatic food webs. And, scientists have developed a screening technique to determine which chemicals pose the greatest risk to the environment.

According to the study led by the U.S. Geological Survey, two traits were identified that indicate how chemicals can build up and reach toxic levels: how easily a [chemical](#) is broken down or metabolized by an organism and the chemical's ability to dissolve in water.

These traits account for how most chemicals concentrate, or biomagnify, in ever-higher levels as one goes up the [food chain](#) from its base to its top predators, such as fish, people, or polar bears. Chemicals that have the ability to biomagnify, such as DDT, can have adverse effects on human and wildlife health and the environment.

"Chemical manufacturers and regulators can use this information to reduce the risks of harmful chemical exposures to ecosystems and the fish, wildlife and people who live in them," said David Walters, a USGS research ecologist and lead author of the study. "By screening for these two characteristics, we can identify chemicals that pose the greatest risk of the thousands that are on the market and for new ones being developed."

The study found that poorly metabolized compounds tend to remain in animal tissues and are passed up the food chain in higher, more toxic amounts as one animal is eaten by another and so on. Likewise, compounds that don't dissolve well in water accumulate in animal fats, ultimately exponentially increasing in top predators.

Beyond these chemical properties, the researchers found that certain ecosystems and food webs are more vulnerable to biomagnification than others. For example, extremely high biomagnification occurred in ocean

food webs that include birds and mammals. The authors noted this may be in part due to longer food chains in these ecosystems that is, many levels and kinds of predators - and because warm-blooded animals need to consume more food than do cold-blooded animals like fish.

Building upon these results, the researchers developed a model of biomagnification based upon how chemicals metabolize and dissolve in water. The likelihood that a chemical would biomagnify was highest - nearly 100 percent—for slowly metabolized compounds such as chlorinated flame retardants and PCBs, or polychlorinated biphenyls, regardless of their solubility in water.

We need to learn from our previous mistakes and have more informed and responsible design and use of chemicals in the environment," said Karen Kidd, a Canada Research Chair at University of New Brunswick Saint John and co-author of the study. "Our global review provides a straightforward approach for reducing the use of chemicals with the properties to concentrate through food webs. This is a critical step for decreasing risks for humans and wildlife from potentially harmful chemical exposures in foods."

Since the emergence of DDT as a global problem for wildlife in the 1950s and 60s, science has kept a close watch on the behavior of persistent organic pollutants, especially chemicals that may concentrate through [food webs](#) to potentially toxic levels in wildlife and humans. Many are resistant to environmental degradation and remain in the environment for decades. While biomagnification can be measured in the laboratory, said Walters, it is best determined by measuring how much the chemical increases with each step in the food chain in wild animal populations.

Provided by US Geological Survey

Citation: Mystery solved: Traits identified for why certain chemicals reach toxic levels in food webs (2016, April 20) retrieved 9 April 2024 from <https://phys.org/news/2016-04-mystery-traits-chemicals-toxic-food.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.