

## Two different collective oscillations of electrons occurring on gold nanoparticles observed for the first time

## April 27 2016



Fig. 1 (a) Diagram of the measurement system to observe the decay process of electron collective motion. The laser beam to the photoemission electron microscope is divided into two, and by delaying the time of the second beam, an image of the electrons emitted from the gold nanoparticles are shown in high resolution, like time-lapse imaging. (b) The photoemission intensity of both dipole and quadrupole plasmon resonance modes as function of the delay time between the pump and probe laser pulses. These results indicate that dipole and quadrupole plasmon resonance exists with different dephasing times. Credit: Hokkaido University



The research group of Professor Hiroaki Misawa of Research Institute for Electronic Science, Hokkaido University and Assistant Professor Atsushi Kubo of the Faculty of Pure and Applied Sciences, University of Tsukuba, have successfully observed the dephasing time of the two different types of collective motions of electrons generated on the surface of a gold nanoparticle for the first time in the world, by combining a laser that emits ultrashort light pulses with a photoemission electron microscope.

When gold is reduced to the size in nanometer scale, its color is red instead of gold. When <u>gold nanoparticles</u> are exposed to light, the collective oscillations of <u>electrons</u> existing on the localized surface of the gold causes red light to be strongly absorbed and dispersed.

This phenomenon is called Surface Plasmon Resonance. The red color of stained glass is also a result of this phenomenon. Recently, gold nanoparticles have been widely used in various fields, such as application in pregnancy tests.

This collective oscillations of electrons on the surface of gold nanoparticles caused by light was considered to be a phenomenon that sustained only for an extremely short time, and difficult to measure due to this shortness.

Our research group developed a methodology to measure the dephasing time of the collective oscillations of electrons occurring on the surface of gold nanoparticles by combining a laser that emits ultrashort light pulses of a few femtoseconds (1 femtosecond =  $10^{-15}$  seconds), and a photoemission electron microscope in <u>high spatial resolution</u>.

When measured by this technique, the different dephasing times of the two different collective oscillations, namely dipole and quadrupole surface plasmon modes, could be resolved and identified as 5



femtoseconds and 9 femtoseconds, respectively.

Research using <u>gold</u> nanoparticles as optical antennae to harvest light for photovoltaic cell and an artificial photosynthesis system that can split water to obtain hydrogen is progressing. The successful measurement of the dephasing time of the collective oscillations of electrons is considered to be a useful guideline in developing these systems.

**More information:** Quan Sun et al. Dissecting the Few-Femtosecond Dephasing Time of Dipole and Quadrupole Modes in Gold Nanoparticles Using Polarized Photoemission Electron Microscopy, *ACS Nano* (2016). DOI: 10.1021/acsnano.6b00715

Provided by Hokkaido University

Citation: Two different collective oscillations of electrons occurring on gold nanoparticles observed for the first time (2016, April 27) retrieved 1 May 2024 from <a href="https://phys.org/news/2016-04-oscillations-electrons-gold-nanoparticles.html">https://phys.org/news/2016-04-oscillations-electrons-gold-nanoparticles.html</a>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.