


# Single-molecule sequencing technology assists microbial DNA methylation study

April 12 2016



Phylogenetic tree of 230 sequenced organisms. Outer bars indicate the number and types of active MTases detected per genome. Credit: Fig. 1A from Blow et al. (2016)

Researchers sequenced 230 diverse archaeal and bacterial genomes to learn more about the roles DNA methylation plays in prokaryotes.

The epigenome of a cell is the collected set of changes made to specific bases in its genomic DNA that affect how the genome is actually used and results from chemical modification (usually methylation.)

Uncovering the breadth of the planet's microbial epigenomic diversity can help researchers better understand the biology of [microbes](#) that play a role in the global nutrient cycles, as well as revealing genomic regulatory systems that could be useful for energy and environment applications.

DNA methylation, the most common epigenetic change, is a process eukaryotes use to regulate [gene expression](#), for example, keeping certain genes from turning on. Though prokaryotes (bacteria and archaea) are also known to have methylated DNA, the roles this process might play in these [single cell organisms](#) is less well understood.

To learn more, a team including researchers at the U.S. Department of Energy Joint Genome Institute (DOE JGI), a DOE Office of Science User Facility, relied on single-molecule, real-time (SMRT) sequencing at the DOE JGI and Pacific Biosciences to reveal DNA methylation patterns in 230 bacterial and archaeal genomes. They found evidence of DNA methylation in 215 microbes (93 percent of those sequenced). These data enabled the annotation of 600 enzymes that methylate DNA (MTases), a massive increase over known annotations. While many DNA methylating enzymes are part of restriction modification systems, consistent with their known role in defense against phages and viruses, the findings suggest that a substantial number of others may be involved in genome regulation, and have a more crucial role in prokaryotic physiology and biology than had been previously suspected.

By mapping and characterizing the epigenetic changes, scientists can associate those targeted genes with environmental adaptations and metabolic activities. Better understanding of such controls on gene expression and under what circumstances they are observed will improve the ability to predict when and where such microbes are detected. In addition, this will inform how microbes interact with plants and other microbes involved in DOE mission interests such as plant bioenergy

feedstock growth, advanced biofuel generation, and soil carbon processing. The report was published February 12, 2016 in *Plos Genetics* and was highlighted February 29, 2016 in Nature Genetics Reviews.

**More information:** Matthew J. Blow et al. The Epigenomic Landscape of Prokaryotes, *PLOS Genetics* (2016). [DOI: 10.1371/journal.pgen.1005854](https://doi.org/10.1371/journal.pgen.1005854)

Provided by DOE/Joint Genome Institute

Citation: Single-molecule sequencing technology assists microbial DNA methylation study (2016, April 12) retrieved 11 May 2024 from <https://phys.org/news/2016-04-single-molecule-sequencing-technology-microbial-dna.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.