

# New tool refines exoplanet search

April 11 2016

---



The tool that allowed the team to improve planet hunting in the near-infrared—a cell that contains methane gas. Credit: Peter Plavchan.

Planet-hunting is an ongoing process that's resulting in the discovery of

more and more planets orbiting distant stars. But as the hunters learn more about the variety among the tremendous number of predicted planets out there, it's important to refine their techniques. New work led by Carnegie's Jonathan Gagné, Caltech's Peter Gao, and Peter Plavchan from Missouri State University reports on a technological upgrade for one method of finding planets or confirming other planetary detections. The result is published by *The Astrophysical Journal*.

One of the most-popular and successful techniques for finding and confirming planets is called the radial velocity method. A planet is obviously influenced by the gravity of the star it orbits; that's what keeps it in orbit. This technique takes advantage of the fact that the planet's gravity also affects the star in return. As a result, astronomers are able to detect the tiny wobbles the planet induces as its gravity tugs on the star. Using this method, astronomers have detected hundreds of exoplanets.

For certain kinds of [low-mass stars](#), however, there are limitations to the standard radial velocity method, which can cause false positives—in other words, find something that looks like a planet, but isn't.

To address this issue, Gagné, Gao, and Plavchan decided to use the [radial velocity technique](#), but they examined a different, longer wavelength of light.

"Switching from the visible spectrum to the near-infrared, the wobble effect caused by an orbiting planet will remain the same regardless of wavelength," Gagné explained. "But looking in the near-infrared will allow us to reject false positives caused by sunspots and other phenomena that will not look the same in near-infrared as they do in visible light,"

Radial velocity work in the near-infrared wavelengths has been conducted before, but it has trailed behind planet hunting in the [visible](#)

[spectrum](#), partially due to technical challenges. The research team was able to develop a better calibration tool to improve the overall technology for near-infrared radial velocity work, which should make it a better option going forward.

They examined 32 low-mass stars using this technological upgrade at the NASA Infrared Telescope Facility atop Mauna Kea, Hawaii. Their findings confirmed several known [planets](#) and binary systems, and also identified a few new planetary candidates.

"Our results indicate that this planet-hunting tool is precise and should be a part of the mix of approaches used by astronomers going forward," Gao said. "It's amazing to think that two decades ago we'd only just confirmed exoplanets actually existed and now we're able to refine and improve those methods for further discoveries."

**More information:** "A High-Precision NIR Survey for RV Variable Low-Mass Stars," Jonathan Gagné et al., 2016, to appear in the *Astrophysical Journal* [arxiv.org/abs/1603.05998](https://arxiv.org/abs/1603.05998)

Provided by Carnegie Institution for Science

Citation: New tool refines exoplanet search (2016, April 11) retrieved 20 April 2024 from <https://phys.org/news/2016-04-tool-refines-exoplanet.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.