


# Bioengineers show why lab-made stem cells might fail: Errors in DNA folding

May 10 2016



When the same linear DNA sequence is divided into different sized segments to analyze its 3-D conformation, different chromatin folding patterns emerge. In this image, the same region around the Sox2 gene is divided into 3, 9, 30, and ~250 bins. The Penn researchers' approach allows us to divide this region into ~250 bins, thus providing a relatively 'high resolution' snapshot of the 3-D conformation of the DNA. Credit: University of Pennsylvania

Induced pluripotent stem cells hold promise for regenerative medicine

because they can, in theory, turn into any type of tissue and because they are made from a patient's own adult cells, guaranteeing compatibility. However, the technique that turns adult cells into these iPS cells is not foolproof; after reverting to their pluripotent state, these cells don't always correctly differentiate back into adult cells.

Researchers from the University of Pennsylvania have now discovered one of the reasons why: the reversion process does not always fully capture the way a cell's genome is folded up inside its nucleus. This folding configuration directly influences gene expression and therefore the functionality of the cell.


The new study shows that current techniques might not produce iPS cells that are equivalent to the pluripotent stem cells found in embryos, as some clones retain folding patterns that partially resemble those found in the adult cells from which they are made.

Led by Jennifer Phillips-Cremins, assistant professor in the School of Engineering and Applied Science's Department of Bioengineering, and Jonathan Beagan, a graduate student in her lab, the study, published in the journal *Cell Stem Cell*, also suggests ways of minimizing these folding errors.

Though techniques for reverting [adult cells](#) into iPS cells have existed for a decade and avoid the issues surrounding the use of [embryonic stem cells](#) that have stymied research into regenerative medicine, clinical investigations of these cells have been cautious and slow. IPS cells can fail to correctly differentiate into the desired tissue. Moreover, there are also concerns that the resulting tissue could have unforeseen genetic abnormalities or could become cancerous.

Even outside the clinical applications, many researchers are interested in iPS cells as a way of generating a "disease in a dish." Rather than taking

a tissue sample from a patient with a genetic disorder, which is especially challenging when the affected organ is the brain, researchers could use iPS cells derived from that patient's [skin cells](#) to grow model organs as needed. Observing the development of those tissues could provide clues to the progression of the disease, as well as serve as ideal test-beds for treatments not yet approved for use in humans.



The 3-D DNA conformation around a critical gene, Sox2, is markedly different between embryonic stem cells, neural progenitor cells, and some induced pluripotent stem cells. These differences are visualized using heatmaps, and illustrations below represent cartoon interpretations of how the DNA may be folding in each cell type based on the heatmap data. Credit: University of Pennsylvania

In both clinical and research applications, however, the traits that allow for the generation of "high quality" iPS cells capable of correctly differentiating into the desired tissue with no genetic abnormalities is unclear.

"We know there is a link between the topology of the genome and gene expression," Phillips-Cremins said, "so this motivated us to explore how the genetic material is reconfigured in three dimensions inside the nucleus during the reprogramming of mature brain cells to pluripotency. We found evidence for sophisticated configurations that differ in important ways between iPS cells and embryonic stem cells."

Phillips-Cremins' area of research is "3-D epigenetics," or the way that the folding of DNA influences gene expression. Classic epigenetic marks are chemical modifications on top of the DNA sequence that provide an additional layer of information on top of the long sequence of base-pair letters. Looking at these marks in a linear fashion does not reveal the whole picture, however, as genome folding can bring two disparate regions of the DNA into spatial and functional contact.

By applying experimental and computational techniques that the Cremins lab has developed, her group was able to identify folding patterns in iPS cells that had previously been unseen.

"Previous methods provided data analogous to an old television, with large, black-and-white pixels," Phillips-Cremins said. "One could generate a blurry image and tell there was a person on the screen, but it would be difficult to parse finer-scale facial features. We employed methods to create high-resolution maps of genome folding, so we could distinguish detailed topological features and evaluate their similarities and differences among traditional embryonic stem cells, iPS cells and mature, [differentiated cells](#)."

The approach Beagan used to create the high-resolution maps involves fixing the DNA such that its 3-D folding patterns are preserved prior to sequencing. Sections of the linear genetic sequence that are separated by marked distances but are spatially adjacent when the DNA is folded are chemically glued together. As a result, two distant parts of the linear sequence will end up in the same string of hybrid DNA and will thus be detected together when the DNA is sequenced.

Analyzing these hybrid pieces provides information that allows the researchers to infer which DNA segments are adjacent to each other in the genome's folded state. Critically, the Cremins lab's approach targets only specific sites in the genome, which allows high-resolution analysis across these regions to be much more easily achievable than with alternative genome-wide approaches. The maps reported by the Penn Engineering lab are the highest-resolution maps of genome folding to date in iPS cells.

Phillips-Cremins and her computational team can plot the sequencing data in heat maps, thereby providing a picture of the DNA segments that are spatially adjacent to each other in the 3-D nucleus of stem cells.

"Each of the pixels on our heat maps," Beagan said, "is a representation of the frequency that any two given segments in the genome interact. You can represent the contact frequency numbers as colors and the entire region of DNA as a heat map. You end up seeing intriguing patterns of high and low intensity, and from these patterns one can infer the folding configurations of the genome."

The Penn researchers targeted several locations along the genome to perform their computational analyses, comparing the iPS cells to the cells from which they were generated and to the embryonic stem cells they ideally should perfectly replicate.

They found that traditional embryonic stem cells and mature, differentiated brain cells had strikingly different genome folding patterns. Surprisingly, however, the genetic material from iPS cells did not fold in a manner that perfectly resembled traditional embryonic stem cells but instead exhibited traces of the 3-D configurations of the [brain cells](#) from which they were derived.

"We found marked differences among the heatmaps we generated for each cell type," Beagan said. "Our observations are important because they suggest that, if we can push the 3-D genome conformation of cells that we are turning into iPS cells to be closer to that of embryonic [stem cells](#), then we can possibly generate iPS cells that match gold-standard [pluripotent stem cells](#) more rapidly and efficiently."

Importantly, the researchers also found they could make iPS cells with more accurate folding configurations by changing the medium in which they are grown. This result provides a window of insight into possible ways that researchers and clinicians could better engineer iPS cells for experiments and treatments. The Phillips-Cremens lab already has experiments underway to engineer 3-D genome folding in stem [cells](#) to control [gene expression](#) on demand.

Provided by University of Pennsylvania

Citation: Bioengineers show why lab-made stem cells might fail: Errors in DNA folding (2016, May 10) retrieved 25 April 2024 from <https://phys.org/news/2016-05-bioengineers-lab-made-stem-cells-errors.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------