

Cancer-fighting properties of horseradish revealed

May 18 2016



Horseradish plant

Horseradish contains cancer-fighting compounds known as glucosinolates. Glucosinolate type and quantity vary depending on size and quality of the horseradish root. For the first time, the activation of cancer-fighting enzymes by glucosinolate products in horseradish has been documented.

The humble [horseradish](#) may not be much to look at, but a recent University of Illinois study shows that it contains compounds that could help detoxify and eliminate cancer-causing free-radicals in the body.

"We knew horseradish had health benefits, but in this study, we were able to link it to the activation of certain detoxifying enzymes for the first time," says U of I crop scientist Mosbah Kushad.

Kushad's research team had previously identified and quantified the compounds responsible for the cancer-fighting compounds, known as glucosinolates, in horseradish, noting that horseradish contains approximately 10 times more glucosinolates than its superfood cousin, broccoli.

"No one is going to eat a pound of horseradish," Kushad points out. Luckily, a teaspoon of the pungent condiment is sufficient to get the benefit.

In the new study, Kushad and his team looked for the [products](#) of glucosinolate hydrolysis, which activate enzymes involved in detoxification of cancer-causing molecules. They compared the quantity and activity of these products in 11 horseradish strains rated U.S. Fancy, U.S. No. 1, or U.S. No. 2. The USDA puts fresh-market horseradish in these categories based on diameter and length of the root.

"There was no information on whether the USDA grade of the horseradish root is associated with cancer preventive activity, so we

wanted to test that," Kushad explains.

The group found that the higher-grade U.S. Fancy accessions had significantly more glucosinolates than U.S. No. 1. Concentrations of various glucosinolate hydrolysis products differed according to USDA grade, with U.S. Fancy having greater allyl isothiocyanate (AITC) and U.S. No. 1 having greater 1-cyano 2,3-epithiopropene (CETP).

The two [compounds](#) differ, with CETP being a comparatively weaker cancer-fighter than AITC. Still, the detection of CETP in horseradish is noteworthy, according to Kushad. "To our knowledge, this is the first detection and measurement of CETP from horseradish," he says.

The team suggests that AITC is a good dietary anti-carcinogen, not only because it activates the enzyme responsible for detoxifying cancer-causing molecules, but also because a large proportion of it, 90 percent, is absorbed when ingested.

More information: Kang-Mo Ku et al. Correlation of Quinone Reductase Activity and Allyl Isothiocyanate Formation Among Different Genotypes and Grades of Horseradish Roots, *Journal of Agricultural and Food Chemistry* (2015). [DOI: 10.1021/jf505591z](https://doi.org/10.1021/jf505591z)

Provided by University of Illinois at Urbana-Champaign

Citation: Cancer-fighting properties of horseradish revealed (2016, May 18) retrieved 24 June 2024 from <https://phys.org/news/2016-05-cancer-fighting-properties-horseradish-revealed.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.