

Study suggests less snow and earlier snowmelt at higher elevations in warming climate

May 26 2016



Sundial Peak, in the Wasatch Mountains, with Lake Blanche (elevation 8920 feet, 2718 meters) in the foreground, May 2016. Credit: David White.

Spring snowpack, relied on by ski resorts and water managers throughout

the Western United States, may be more vulnerable to a warming climate in coming decades, according to a new University of Utah study.

The study, accepted for publication in *Geophysical Research Letters*, models the year-to-year variability in precipitation and temperature in Utah's Wasatch Mountains and other ranges in the West. Jason Scalzitti, a graduate student in atmospheric sciences, and professors Court Strong and Adam Kochanski found that above a threshold elevation, the amount of spring snowpack is dependent more on the amount of precipitation in a year than the temperature. In other words, whether a year is wet matters more than if it's warm. But below that threshold, temperature matters more. By the end of the century, according to the study, that threshold will move uphill by around 800 feet in the Wasatch and more in the Sierra Nevada, Cascades and parts of the Rocky Mountains.

"In the past we've thought mainly about total precipitation as an indicator of how good the ski season's going to be," says Strong. "As we move into the future, especially at elevations below the threshold, temperature increases in importance."

Zooming in on the Wasatch

Strong and his colleagues based their work on a high-resolution, regional-scale climate model called the Weather Research and Forecasting (WRF) model. Conventional climate models make calculations on grids 62 miles (100 km) on a side. For reference, less than 20 miles separate Salt Lake City, west of the Wasatch, and Park City, Utah, on the east. "You can't even see the Wasatch Range at that resolution," Strong says. Although the coarser grid of global climate models works well in flat topography, such as the Great Plains, the complex terrain of the Intermountain West requires a finer resolution.

The researchers employed a technique called "dynamical downscaling,"

telescoping the model grid into smaller and smaller grid cells in order to capture fine-scale atmospheric processes affecting local climate. In the team's final simulations, the Wasatch Range was modeled at a resolution of about 2.5 miles (4 km) to realistically capture impacts of the range's slopes, canyons, and peaks on the local precipitation pattern. They further accounted for future temperature changes in the Great Salt Lake and evaporation from urban irrigation, both of which contribute moisture to the air. The team projected regional future climate forward to the year 2100 using a business-as-usual carbon emissions scenario that assumes greenhouse gas emissions will continue to increase at the same rate as today.

Crossing the threshold

They found that at high elevations, temperatures remain cold enough throughout the spring to turn precipitation to snow and to keep the snow on the ground from melting. Amount of precipitation is the main factor determining how much snow is on the ground during the critical spring months. Going down toward the valley floors, however, temperatures rise, and even in years with high precipitation, the slopes experience more rain and more melting. Temperature becomes the primary driver of the depth of spring snowpack.

The threshold elevation between precipitation-controlled and temperature-controlled snowpack, currently sits at about 6500 feet (1980 meters) in the Wasatch, near the base elevation of the ski resorts in Park City. In future projections, however, the threshold elevation rises to 7300 feet (2230 meters).

In the simulations, modeled threshold elevations rose in mountain ranges all over the West - by about 800 feet (250 meters) in Colorado's Rockies, about 980 feet (300 meters) in California's Sierra Nevada and Washington's Cascades and more than 1400 feet (432 meters) in the

middle Rockies of Idaho and Wyoming.

Impacts in peaks and valleys

Four Wasatch ski resorts, Solitude, Snowbird, Alta, and Brighton, sit well above the end-of-century 7300 foot threshold. But the rest, including venues from the 2002 Winter Olympics, sit at base elevations between 5500 and 7200 feet. The peak elevations of all resorts except one, Nordic Valley, extend up above the 7300 foot future threshold.

An overall warming trend doesn't mean that every year will be a bad year for low-elevation resorts. But below the threshold elevation, resorts will be more susceptible to warm years. "Let's say we get the same amount of storms every year," says study co-author Kochanski. "Above the threshold, the resorts will probably be fine. For the others, even if we have the same [precipitation](#), they may be in trouble because they could get more rain instead of snow, and the snowpack will diminish faster."

Although impacts to ski resorts could affect Utah's economy, another implication of diminished snowpack affects nearly all Utahns - shrinking water resources. Melting spring snowpack fills reservoirs, providing water for the residents of the Salt Lake Valley. Spring snowpack amounts are a key indicator for water managers of how much water they'll have available in reservoirs for the coming year.

"They look at that as how much water is available in the form of snow to melt and capture in the reservoirs," Strong says. "That will be down in the future. Even if we have the same amount of water coming into the system, it will be melting earlier and faster. If we want to supply that to a growing population, then we need increased storage capacity."

Strong and Kochanski say they can now use their model as a framework to continue exploring questions about future climate and snowpack

variation, both by exploring the impacts of different carbon emissions scenarios and by looking at mountain ranges around the world, such as the Himalayas.

More information: Jason Scalzitti et al, Climate change impact on the roles of temperature and precipitation in western U.S. snowpack variability, *Geophysical Research Letters* (2016). DOI: [10.1002/2016GL068798](https://doi.org/10.1002/2016GL068798)

Provided by University of Utah

Citation: Study suggests less snow and earlier snowmelt at higher elevations in warming climate (2016, May 26) retrieved 26 April 2024 from <https://phys.org/news/2016-05-earlier-snowmelt-higher-elevations-climate.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.