

## Electronic micro labs control chemical processes from the inside

May 3 2016



PhD student Abhishek Sharma performs one of many micro lab production steps. He positions electrodes in order to coat a part of the wafer. Credit: RUB, Damian Gorczany

Chemists at the Ruhr-Universität Bochum, together with their project



partners, have developed tiny electronic components that can control chemical processes from the inside. These micro labs are  $140 \times 140 \times 60$  micrometres in size. They can sense their environment and control chemistry via voltage signals. One day, they shall acquire the capability to evolve.

Autonomous micro labs the size of cells that monitor chemical systems from the inside – pure fiction, as yet. But several crucial steps have been taken.

Lablet, is the name Prof Dr John McCaskill from the Ruhr-Universität Bochum has given to the microscopic electronic component that he has been developing in the recent years, together with numerous partners. As researcher at the Faculty of Chemistry and Biochemistry, he heads the EU project "Microscopic Chemically Reactive Electronic Agents".

The international team aims at electronically controlling <u>chemical</u> <u>reactions</u> from the inside, rather than aiding them via large interface structures from the outside. "Just imagine what a chemist could achieve if he was as small as a cell," says McCaskill.

## **Electrodes control chemistry via voltage signals**

Developed in the course of the project, the tiny labs, also called lablets, are  $140 \times 140 \times 60$  micrometres in size. They have the capacity of controlling various chemical reactions in aqueous solutions and absorbing information from their environment. To this end, they are equipped with electronics and microscopic electrodes.





Together with his team, John McCaskill develops electronic micro-agents that control chemical processes. Credit: RUB, Damian Gorczany

A voltage applied to the electrode can supply the energy required for initiating a <u>chemical</u> reaction. Different voltages cause different reactions – which ones depends on the previous history and, for example, the molecules available in the environment.

The researchers have developed a supercapacitor for the micro labs that currently provides energy for autonomous operation for about 20 minutes. For recharging and programming, the lablets have to bind to a smart surface.



## Micro labs designed with the capacity to evolve

When designing the lablets, the chemists have taken several cues from nature. They are set up in such a way that they can transfer their programming to other units by passing along a sequence of voltage signals. One day, the lablets should be able to pair up autonomously and acquire the capability to evolve.

The project team is now working at full stretch at integrating all functions developed so far in one single micro lab.

Provided by Ruhr-Universitaet-Bochum

Citation: Electronic micro labs control chemical processes from the inside (2016, May 3) retrieved 3 May 2024 from <u>https://phys.org/news/2016-05-electronic-micro-labs-chemical.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.