

Water on the moon?

May 30 2016



This illustration shows a promising new technique for locating water on the Moon. Galactic cosmic rays (GCR) that penetrate the lunar surface, when encountering a layer with material containing hydrogen atoms (like water), trigger the ejection of protons (red spheres) that can be detected by the appropriate instrumentation in an orbiting satellite. Credit: Schwadron et al. 2016

Prior to the Apollo missions to the moon, scientists speculated that



volatiles - including water - may have accumulated in permanently shaded regions at the poles. Then the Apollo era brought the return of lunar samples, enabling real measurements: They found none of the water-bearing minerals common on Earth. Over the past ten years, however, several developments have reinvigorated the discussion. In particular, new analyses of volcanic glasses in the sample returns have inferred the presence of water in the moon's interior. Meanwhile, several new lunar missions have been launched. The ones using neutron spectroscopy to search for water have come up with mixed conclusions, but those using infrared spectroscopy seemed to reach unambiguous identification of water on the lunar surface, although in disagreement with the neutron experiments.

CfA astronomers Anthony Case and Justin Kasper were members of a team of astronomers who propose a new method to detect hydrated material on the moon - like water - by measuring the strength of protons coming from the lunar surface with the CRaTer instrument (Cosmic Ray Telescope for the Effects of Radiation) on the Lunar Reconnaissance Orbiter, a NASA robotic orbiter launched in 2009. Cosmic rays from the galaxy, when striking the lunar surface, will knock protons out of material on the surface which can be detected by the CRaTER instrument. The team completed a set of laboratory tests using high energy particle accelerators to simulate the effects of <u>cosmic rays</u> on materials containing hydrogen, and found that the presence of hydrogen - in water for example – actually suppresses the overall proton emission. The implication is that if water is present near the poles of the moon, a scan across the <u>lunar surface</u> should show a clear reduction in numbers of protons as it crosses the poles.

Actually, the CRaTER scans found an increase in the proton emission at the poles. The scientists soon realized that there were some effects, originally thought to be negligible, that were responsible. Protons and neutrons, released from material below the surface down to about ten



centimeters, will collide with other atoms and produce the emission of secondary particles. The enhancement of these secondary <u>protons</u> is indeed consistent with the presence of hydrogen. But it turns out that there are other possible solutions as well, and the team is continuing to investigate them. Meanwhile their current paper shows that the technique of using CRaTER measurements to search for water is at least in principle possible, and when the remaining issues are resolved, the techniques could be used in other missions to probe other solar system bodies.

More information: N.A. Schwadron et al. Signatures of volatiles in the lunar proton albedo, *Icarus* (2016). <u>DOI:</u> 10.1016/j.jcarus.2015.12.003

Provided by Harvard University

Citation: Water on the moon? (2016, May 30) retrieved 3 May 2024 from <u>https://phys.org/news/2016-05-moon.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.