

Research finds skull condition thought extinct is actually widespread

May 17 2016, by Ann Ross



Example of a skull exhibiting cribra orbitalia—note the porous areas in the top of the eye sockets. Credit: Ann Ross

Some forensic anthropologists thought the skull condition called cribra orbitalia (CO) was a thing of the past – but new research from North Carolina State University and the University of the Witwatersrand finds that it not only still exists, but is fairly common in both North America and South Africa.

CO is a condition in which the bone inside the eye sockets becomes porous. It is not known to cause any [adverse health effects](#), but is generally regarded as being caused by [iron deficiency anemia](#).

The condition has traditionally been used by anthropologists to assess diet and health in prehistoric populations. For example, the presence of CO could tell researchers that a population was not getting a sufficiently varied diet.

"But there's been a lot of debate about the prevalence of CO in modern populations, with some saying it had effectively disappeared," says Ann Ross, director of the Forensic Sciences Institute at NC State and co-author of a paper on the work. "We wanted to know if CO was still extant and, if so, how common it is in modern populations, relative to earlier eras."

For this study, the researchers looked at modern, historic and prehistoric human remains from South Africa, North Carolina and the Western Hemisphere Database. Altogether, the researchers evaluated data on 844 [skulls](#): 245 prehistoric, 381 historic (as recent as the early 20th century) and 218 modern.

Their findings were surprising.

The researchers found that CO was not only present in modern populations, but that it was not even uncommon.

For example, the researchers found that two of the five modern North American juvenile skulls evaluated in the study – 40 percent – had CO. And 15 of the 60 South African juveniles evaluated in the study – 25 percent – had CO.

"We thought we might see some CO, but not to the extent that we did,"

Ross says. "The high rates may stem from the fact that these remains were part of forensic cases – there were often related to cases of homicide or neglect. These cases are not representative of health for all children."

Overall, the researchers found that 12.35 percent of modern North Americans and 16.8 percent of modern South Africans, across all age groups, had CO.

Both rates are higher than their historic counterparts. Only 2.23 percent of historic South African skulls evaluated had CO, and only 6.25 percent of historic North American skulls. Even the prehistoric North American skulls had a lower rate of CO, at 11.86 percent.

"We think the increased prevalence of CO in the modern skulls may be due to intestinal parasites in some populations and iron-poor diet," Ross says.

"These findings drive home the fact that disadvantaged socioeconomic groups, and parts of the developing world, are still struggling with access to adequate nutrition," Ross adds. "Corn may give people a full belly, but it's not going to give people all of the nutrients they need to be healthy."

More information: Maryna Steyn et al. Cribra Orbitalia: Prevalence in Contemporary Populations, *Clinical Anatomy* (2016). [DOI: 10.1002/ca.22734](https://doi.org/10.1002/ca.22734)

Provided by North Carolina State University

Citation: Research finds skull condition thought extinct is actually widespread (2016, May 17) retrieved 25 April 2024 from

<https://phys.org/news/2016-05-skull-condition-thought-extinct-widespread.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.