

# Sweet spot found for foraging manta rays

May 5 2016

---



A team of international scientists has unlocked the foraging secrets of manta rays at Lady Elliot Island, which could help conservation efforts for the vulnerable species.

Led by The University of Queensland, the team found manta rays only

feed when the biomass of their food source—tiny creatures known as zooplankton—passes a critical threshold, which happens two hours before high tide.

"This is the first study of the feeding and prey availability at any site where reef manta rays are known to aggregate," said paper co-author Professor Anthony Richardson of UQ's School of Mathematics and Physics, and CSIRO Oceans and Atmosphere.

Lady Elliot Island is a well-known gathering site for the mysterious reef manta rays (*Manta alfredi*) in the southern Great Barrier Reef.

"Despite the popular interest worldwide for manta rays, very little is known about their biology and ecology," Professor Richardson said.

"However, further knowledge is essential for conservation."

The researchers visited three sites near Lady Elliott Island to collect samples of prey, and combined this with long-term sighting records for [manta rays](#) in the area.

They performed laboratory and complex statistical analyses, factoring in oceanographic influences, to reach their conclusions.



Study leader Asia Armstrong, from UQ's School of Biomedical Sciences and Project Manta, said manta ray [species](#) were listed as vulnerable to extinction on the International Union for the Conservation of Nature's Red List of Threatened Species.

"Over the past decade, there has been considerable research effort to understand the basic ecology and biology of these animals," she said.

"A remaining gap, however, has been an understanding of their movement ecology in relation to foraging.



"Conserving highly mobile species can be particularly challenging. However analysis of species critical habitats and drivers for their migrations can help inform conservation planning."

The research is published today in *PLOS One*.



**More information:** Asia O. Armstrong et al. Prey Density Threshold and Tidal Influence on Reef Manta Ray Foraging at an Aggregation Site on the Great Barrier Reef, *PLOS ONE* (2016). [DOI: 10.1371/journal.pone.0153393](https://doi.org/10.1371/journal.pone.0153393)

Provided by University of Queensland

Citation: Sweet spot found for foraging manta rays (2016, May 5) retrieved 6 May 2024 from

<https://phys.org/news/2016-05-sweet-foraging-manta-rays.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.